

Determination of optimum feeding rate for growth of Caspian carp, *Cyprinus carpio* (Linnaeus, 1758) fingerlings

Majid Mohammad Nejad Shamoushaki, Zia Khari, Zeynab Eslami

Department of Fishery, Bandar Gaz Branch, Islamic Azad University, Bandar Gaz, Iran.
Corresponding author: M. Mohammad Nejad Shamoushaki,
majid_m_sh@bandargaziau.ac.ir

Abstract. One of the important issues in aquaculture, need to achieve a balance between growth rate and optimal use of fish meal is provided. This study was conducted for 6 weeks in the Sijual Bony Fishes Reproduce and Cultivate Center (Gorgan, Golestan, Iran). Experiments were done 4 treatments and 3 replicates as follows: treatment A = 2.5 % body weight per day, treatment B = 5 % body weight per day, treatment C = 7.5 % body weight per day and treatment D = 10 % body weight per day. Aim to this purpose Caspian carp fingerlings with an average weight of 1.5 ± 0.145 g and average length of 2.97 ± 0.154 cm were distributed in 12 fiberglass tanks (30 number fish per tank) and were cultured for 6 weeks in the same conditions and fed with SFC diet (content: 8.7 % moisture, 11.2 % ash, 32 % protein and 10.5 % fat) four times a day. Results showed that hadn't significant effects in Specific Growth Rate (SGR), Body Weight Index (BWI %), Growth Rate (GR) and Condition Factor (CF) ($P < 0.05$) but hadn't significant effects in Feed Conversion Ratio (FCR) and survival ($P > 0.05$). The results show that there are significant differences with regard to the amount of body weight and body length of fingerlings in different treatments ($P < 0.05$), so that the maximum body weight and body length of carp fingerlings is 7.5 % body weight per day.

Key Words: feeding rate, growth, survival, *Cyprinus carpio*.

Introduction. Fish is a primary source of protein for more than one billion people around the world (Rameshguru et al 2011). Understanding of natural foods and dietary habits of fish culture could be an important factor in providing effective method of nutrition. Although intensive fish culture adaptability of the species with different feeding methods have been proven, but the choice of methods to provide food and nutrition in aquaculture should be considered dietary at patterns of normal behavior (Mohammad Nejad Shamoushaki 2012). One of the most significant attainments in aquaculture is fish feeding due to the high cost of feeding (Aydin et al 2011). Feeding rate, water temperature, and fish size have a significant effect on the fish growth; in particular, the optimum feeding rate has to be found in order to achieve a successful aquaculture operation (Yuan et al 2010). Optimum feeding rate differ according to fish species, size, and rearing system (Cho et al 2003). In addition, feeding rate affects the usage of the nutrients in the feed (Mihelakakis et al 2002). Knowledge of the optimum feeding rate is important to improve best feed efficiency resulting better growth and production (Cho et al 2003; Eroldogan et al 2004). Moreover, knowing of the optimum feeding rate helps to prevent water quality degradation which overfeeding causes (Ng et al 2000; Mihelakakis et al 2002; Webster et al 2002; Eroldogan et al 2004). Feeding rate is especially important for larval fish since they are vulnerable to over- and under-feeding. Therefore, over- or under-feeding causes an increase in disease and mortality (Deng et al 2003). Feeding costs contributes up to 60% of the variable costs of culture systems. Therefore, it is essential to provide a proper and applicable feeding management program. In fact, an inadequate food supply directly affects production costs and water quality (Silva et al 2007; Mihelakakis et al 2002; Ng et al 2000). Overfeeding mostly causes feed spillage,

decreasing in feed efficiency, and polluting the environment. Similarly, underfeeding results in decreased feed efficiency as well as degraded growth (Dediu et al 2011). Therefore, it is essential, in terms of both economy and biology, to determine the optimum feeding rate for growth (Aydin et al 2011). It should be noted that optimal feeding rate is essential not only because of promoting best growth and minimizing feed conversion rate (FCR), but also for economic and environmental aspects, preventing water quality degradation (Yuan et al 2010).

Caspian carp, *Cyprinus carpio* belongs to *Cyprinidae* is one of the most economically important and valuable teleostei in the Caspian Sea. Currently, there is no published information on the effects of feeding rate on growth of Caspian carp fingerlings. It is necessary to have a better understanding of the optimum feeding rate of this species. The objective of the this study was to determine optimum and maintenance feeding rate and the effects of the feeding rate on growth performances and survival for Caspian carp, *C. carpio* fingerlings.

Materials and Methods. This study has been carried out in Sijual Bony Fishes Reproduce and Cultivate Center (Gorgan, Golestan, Iran) in 2011 summer. Initial body weight and length average were 1.5 ± 0.145 g and 2.97 ± 0.154 cm. Given the importance of the physical and chemical performances and their impact on water supply and ultimately the fish growth, these performances were so controlled through the experiment that the amount of dissolved oxygen was fixed on 5.5 - 6 ppm, the temperature $26 \pm 2^{\circ}\text{C}$ and pH 7.5 to 8. Fish were fed SFC (starting food carp) commercial food during (6 weeks) of the study. The feed was administered only during daylight hours. Pertinent characteristics of this feed were: 32% crude protein; 10.5% crude fat; 11.2% ash and 8.7% moisture. This study was operated as long as 6 weeks and in 12 tanks with 30 numbers fish in each tank, with 4 treatments and 3 replicate as: treatment A = 2.5 % body weight per day — BW d^{-1} , treatment B = 5 % body weight per day — BW d^{-1} , treatment C = 7.5 % body weight per day — BW d^{-1} and treatment D = 10 % body weight per day — BW d^{-1} . Daily treatments were fed four times a day (equal meals at 08:00, 11:00, 14:00 and 17:00 h) by hand according to feeding rates, making sure that no feed was left uneaten. Fish were sampled every 2 weeks to evaluate growth in weight and length, for this purpose 10 numbers of fish in each tank were captured, weighed and measured. After each sampling period the amount of feed given was adjusted according to mean weight in each aquarium. From results of the last sample fish performances were evaluated in terms of Feed Conversion Ratio (FCR), Specific Growth Rate (SGR, $\% \text{ d}^{-1}$), Body Weight Index (BWI %), Growth Rate (GR, g d^{-1}), Condition Factor (CF, g/Cm) and Survival (%). These performance indices were calculated as follows (Hung et al 1989; Ronyai et al 1990; Biswas et al 2010):

- $\text{FCR} = \text{total feed intake} / \text{total biomass gain}$
- $\text{SGR} = [(\ln \text{ final weight} - \ln \text{ initial weight}) / \text{rearing duration in days}] \times 100$
- $\text{BWI} = [(\text{body weight final} - \text{body weight initial}) / \text{body weight initial}] \times 100$
- $\text{GR} = (\text{body weight final} - \text{body weight initial}) / \text{rearing duration in days}$
- $\text{BWI} = [(\text{body weight} / \text{total length}^3)] \times 100$
- $\text{Survival} = (\text{number of fish harvested} / \text{number of fish stocked}) \times 100$

SPSS version 13 and a software program for drawing graphs of Excel 2003 have been used to analys all data. All data were analyzed with one-way analyses of variance (ANOVA) and significant means were subjected to a multiple comparison test (Duncan) at $P < 0.05$. When the normality of data did not present, the nonparametric test Kruskal-Wallis to compare treatments and test Mann - Whitney for paired comparison between treatments were used (Mohammad Nejad Shamoushaki 2012).

Results and Discussion. Final body weight and body length of *C. carpio* in different treatments are shown in Table 1. Obtained results at the end of this study showed that feeding rates had significant effects on body weight and body length of *C. carpio* ($P < 0.05$). Also, results show that maximum body weight and body length were in 7.5% BW d^{-1} .

Table 1

The average of body weight and body length of *C. carpio* in different treatments

Treatments	Initial weight (g)	Initial length (cm)	Final weight (g)	Final length (cm)
A 2.5 % BW d ⁻¹	1.5 ± 0.145	2.97 ± 0.154	2.046 ± 0.076 ^a	3.327 ± 0.097 ^a
B 5 % BW d ⁻¹	1.5 ± 0.145	2.97 ± 0.154	2.875 ± 0.212 ^b	3.34 ± 0.566 ^a
C 7.5 % BW d ⁻¹	1.5 ± 0.145	2.97 ± 0.154	3.337 ± 0.085 ^c	4.357 ± 0.365 ^b
D 10 % BW d ⁻¹	1.5 ± 0.145	2.97 ± 0.154	3.14 ± 0.07 ^d	4.297 ± 0.091 ^c

*The small Latin letters show that there are significant differences among different treatments

Comparison average of different feeding rate effects on *C. carpio* growth performances during culture period are shown in Table 2. Results showed that hadn't significant effects in BWI, SGR, GR and CF (P<0.05) but hadn't significant effects in FCR and survival (P>0.05).

Table 2

Effects of different feeding rate on growth performances in *C. carpio*

Indicators	Treatment A	Treatment B	Treatment C	Treatment D
FCR	4.65 ± 2.438 ^a	5.106 ± 1.768 ^a	7.737 ± .205 ^a	7.465 ± 0.375 ^a
SGR (% d ⁻¹)	7.24 ± 3.45 ^a	13.83 ± 2.85 ^b	14.64 ± 0.19 ^c	16.37 ± 0.61 ^d
BWI (%)	7.78 ± 3.18 ^a	14.96 ± 7.43 ^b	13.17 ± 0.31 ^b	18.79 ± 2.08 ^c
GR (g d ⁻¹)	0.011 ± 0.004 ^a	0.026 ± 0.001 ^b	0.0277 ± 0.0007 ^b	0.035 ± 0.003 ^c
CF (g/cm)	1.36 ± 0.12 ^a	1.94 ± 0.44 ^b	1.32 ± 0.24 ^a	1.26 ± 0.046 ^c
Survival (%)	100 ^a	100 ^a	100 ^a	100 ^a

*The small Latin letters show that there are significant differences among different treatments

Fish requires a daily ration for maximum growth (Chua & Teng 1982; Kaiser et al 2012). Fish feeding is one of the most important performances in aquaculture because of high feed costs. Overfeeding causes degradation of water quality; consequently, the fish growth reduces and cost increases. On the other hand, underfeeding is undesirable. Therefore, it is critical to find the optimum feeding rate from both economical and biological point of view (Aydin et al 2011). In this work, maximum growth of Caspian carp fed at 7.5% BW d⁻¹. The optimal feeding strategies augment growth performance, survival, and food conversion ratios. Additionally, it contributes to minimize food wastage, reduce size variation, and accordingly, increase production efficiency (Goddard 1996; Kubitzka & Lovshin 1999). In his study, significant differences in BWI, SGR, GR and CF (P<0.05) were observed among treatments (see Table 2). The SGR and GR increased with increasing feeding rate to 10% BW d⁻¹ (P<0.05) but best CF were observed in 7.5% BW d⁻¹ (P<0.05). The highest growth in the present study typically occurred in the 7.5% BW d⁻¹ group. A similar result was also found in sea bass (Russell et al 1996) and *Rachycentron canadum* (Sun et al 2006). The best growth found at the 7.5% and 7% BW d⁻¹.

The quantity and quality of the ration have a significant effect on growth rate, efficiency of feed utilization and chemical composition (Juell & Lekang 2001; Abdelghany & Ahmad 2002; Rowland et al 2005; Bureau et al 2006; Yuan et al 2010). Therefore, it is necessary to determine the optimal feeding rate for different species. Results of other studies are different in various species. Studies with several fish species have revealed that with increasing feeding rate, the growth increases at higher ration levels and decreases at lower ration levels (De Silva & Anderson 1995; Hung & Lutes 1987; Hung 1991; Xiao-Jun & Ruyung 1992; Adebayo et al 2000; Ng et al 2000; Mihelakakis et al 2002; Rosenthal 2000; Tsevis et al 1992). The discrepancies between the results of this study and the findings of others could be explained by size and/or age differences of fish

and experimental conditions such as feeding regimes (Brett 1979; Webster et al 2002; Fiogbe & Kestemont 2003). For instance, Hung et al (1993) observed that optimum feeding rate for striped bass fingerlings (initial body weight 38 g) was 1.0–1.5% BW d⁻¹. In similar size, optimum feeding rate was suggested to be 2.0 BW d⁻¹ for the same species (>27 g) kept by Piper et al (1982). It is well known that feeding rate and feed consumption in relation to body weight decreases as fish grow (NRC 1993; Ng et al 2000; Webster et al 2002). A feeding rate of 4.0% BW d⁻¹ for gilthead sea bream at 3g initial average weight but 6.0% BW d⁻¹ at 1g initial average weight was suggested to be optimal (Kalogeropoulos et al 1992; Alexis et al 1999; Eroldogan et al 2004). Also, another study, Silva et al (2007), showed that increasing feeding rate on *Colossoma macropomum* with 10% BW d⁻¹ results in more growth. Also, studies on other fish such as *Mystus nemurus* (Ng et al 2000), *Dicentrarchus labrax* (Eroldogan et al 2004), *Ictalurus punctatus* (Robinson & Li 1999) and *Piaractus mesopotamicus* (Borghetti & Canzi 1993) showed that feeding rate with 10% BW d⁻¹ fish growth was better.

Conclusions. The result of this research on the effects of the fish feeding rate has been different from other studies. However, research results show that the feeding rates and growth rate are different in different species. Based on growth performance, the highest growth was obtained with fish fed 7.5% BW day⁻¹, in the present study. It appears that, feeding 7.5% BW day⁻¹ to satiation may be accepted as sufficient for on growing of Caspian carp under the conditions of this experiment.

Acknowledgements. This research was supported by the Sijual Bony Fishes Reproduce and Cultivate Center (Gorgan, Golestan, Iran). I also thank Mr. Jabbare for supplying the Caspian carp fingerlings used in this study. I also thank Mr. Maleki, Mr. Shakiba and Mr. Eri for their valuable help during the experiments.

References

- Abdelghany A. E., Ahmad M. H., 2002 Effects of feeding rates on growth and production of Nile Tilapia, common carp and silver carp polycultured in fertilized ponds. *Aquac Res* 33: 415-423.
- Adebayo O. T., Balogun A. M., Fagbenro O. A., 2000 Effect of feeding rates on growth, body composition and economic performance of juvenile clariid catfish hybrid (*Clarias gariepinus* x *Heterobranchus bidorsalis*). *J Aquacult Trop* 15: 109-117.
- Alexis M. N., Nengas I., Fountoulaki E., Papoutsi E., Andriopolou A., Koutsodimou M., Gaubaudan J., 1999 Tissue ascorbic acid levels in European sea bass (*Dicentrarchus labrax*) and gilthead sea bream (*Sparus aurata*, L.) fingerlings fed diet containing different forms of ascorbic acid. *Aquaculture* 179: 447–456.
- Aydın I., Küçük E., Şahin T., Kolotoğlu L., 2011 The effect of feeding frequency and feeding rate on growth performance of juvenile black sea turbot (*Psetta maxima*, Linnaeus, 1758). *J FisheriesSciences.com* 5(1): 35-42.
- Biswas G., Thirunavukkarasu A. R., Sundaray J. K., Kailasam M., 2010 Optimization of feeding frequency of Asian seabass (*Lates calcarifer*) fry reared in net cages under brackishwater environment. *Aquaculture* 305: 26–31.
- Borghetti J. R., Canzi C., 1993 The effect of water temperature and feeding rate on the growth rate of pacu (*Piaractus mesopotamicus*) raised in cages. *Aquaculture* 114: 93–101.
- Brett J. R., 1979. Environmental performances and growth. Bioenergetic and growth. Fish physiology, W. S. Hoar D.J. Randole and J.R Brett, Eds. Vol. 8, Academic Press, New York, USA. Pp.599-675.
- Bureau D. P., Hua K., Cho C. Y., 2006 Effect of feeding level on growth and nutrient deposition in rainbow trout (*Oncorhynchus mykiss* Walbaum) growing from 150 to 600g. *Aquac Res* 37: 1090-1098.
- Cho S. H., Lim Y. S., Lee J. H., Park S., 2003 Effect of feeding rate and feeding frequency on survival, growth, and body composition of ayu post-larvae *Plecoglossus altivelis*. *J World Aquac Soc* 34: 85–91.

- Chua T. E., Teng S. K., 1982 Effects of food ration on growth, condition factor, food conversion efficiency, and net yield of estuary grouper, *Epinephelus salmoides* Maxwell, cultured in floating net cages. *Aquaculture* 27: 273–283.
- Dediu L., Cristea V., Mocanu M., Dicu D., Angelica Docan A., Grecu I., 2011 The effect of feeding frequency on growth performance of rainbow trout fingerlings reared in recirculating system. *AAFL Bioflux* 4(2): 141-145.
- Deng D. F., Koshio S., Yokoyama S., Bai S. C., Shao Q., Cui Y., Hung S. S. O., 2003 Effects of feeding rate on growth performance of white sturgeon (*Acipenser transmontanus*) larvae. *Aquaculture* 217: 589-598.
- De Silva S. S., Anderson T. A., 1995 *Fish Nutrition in Aquaculture*. Aquaculture Series, Chapman & Hall, London.
- Eroldogan O. T., Kumlu M., Aktas M., 2004 Optimum feeding rates for European sea bass (*Dicentrarchus labrax* L.) reared in seawater and freshwater. *Aquaculture* 231: 501–515.
- Fiogbe E. D., Kestemont P., 2003 Optimum ration for Eurasian perch *Perca fluviatilis* L. reared at its optimum growing temperature. *Aquaculture* 216: 243–252.
- Goddard S., 1996 *Feed Management in Intensive Aquaculture*, Chapman & Hall, New York, 194 pp.
- Hung S. S. O., Lutes P. B., 1987 Optimum feeding rate of hatchery-produced juvenile white sturgeon (*Acipenser transmontanus*): at 20 °C. *Aquaculture* 65: 307-317.
- Hung S. S. O., Lutes P. B., Storebakken T., 1989 Growth and feed efficiency of white sturgeon (*Acipenser transmontanus*) sub yearling at different feeding rates. *Aquaculture* 80: 147-153.
- Hung S. S. O., 1991 Nutrition and feeding of hatchery-produced juvenile white sturgeon (*Acipenser transmontanus*): an overview. In: P. Williot (Ed.), proceedings of the First International Symposium on the Sturgeon, Cemagref, Bordeaux, France: 65-77.
- Hung S. S. O., Conte F. S., Hallen E. K., 1993 Effects of feeding rates on growth, body composition and nutrient metabolism in striped bass (*Morone saxatilis*) fingerlings. *Aquaculture* 112: 349–361.
- Juell J. E., Lekang O. I., 2001 The effect of feed supply rate on growth of juvenile perch (*Perca fluviatilis*). *Aquac Res* 32: 459-464.
- Kaiser H., Collett P. D., Vine N. G., 2012 The effect of feeding regimen on growth, food conversion ratio and size variation in juvenile dusky kob *Argyrosomus japonicus* (Teleostei: *Sciaenidae*). *African Journal of Aquatic Science* 36(1): 83–88.
- Kalogeropoulos N., Alexis M. N., Henderson R. J., 1992 Effects of dietary soybean and cod-liver oil levels on growth and body composition of gilthead bream (*Sparus aurata*). *Aquaculture* 104: 293–308.
- Kubitza F., Lovshin L. L., 1999 Formulated diets, feeding strategies, and cannibalism control during intensive culture of juvenile fishes. *Reviews in Fisheries Science* 7: 1–22.
- Mihelakakis A., Tsolkas C., Yoshimatsu T., 2002 Optimization of feeding rate for hatchery-produced juvenile gilthead sea bream *Sparus aurata*. *J World Aquac Soc* 33: 169–175.
- Mohammad Nejad Shamoushaki M., 2012 Effect of feeding frequency on growth performances and survival of *Rutilus rutilus caspicus*. *Journal of Research in Biology* 2(3): 184-189.
- Ng W. K., Lu K. S., Hashim R., Ali A., 2000 Effects of feeding rate on growth, feed utilization and body composition of tropical bagrid catfish. *Aquac Int* 8: 19–29.
- NRC (National Research Council), 1993 *Nutrient Requirement of Fish*. National Academy Press, Washington, DC, USA.
- Piper R. G., McElwain I. B., Orme L. E., McCraren J. P., Fowler L. G., Leonard J. R., 1982 *Fish Hatchery Management. Nutrition and Feeding*, Vol. 4. United States Department of the Interior Fish and Wildlife Service, Washington, DC, USA, p. 254.
- Robinson E. H., Li M. H., 1999 Effect of dietary protein concentration and feeding rate on weight gain, feed efficiency, and body composition of pond-raised channel catfish (*Ictalurus punctatus*). *J World Aquac Soc* 30: 311–318.

- Rameshguru G., Senthilkumar P., Govindarajan B., 2011 Vermiwash mixed diet effect on growth of *Oreochromis mossambicus* (Tilapia). *Journal of Research in Biology* 5: 335-340.
- Ronyai A., Peteri A., Radics F., 1990 Cross breeding of starlet and Lena river sturgeon. *Aquaculture* 6: 13-18.
- Rosenthal A., 2000 Status and prospects of sturgeon farming in Europe. Institute für Meereskunde Kiel Dusternbrooker Weg 20 2300 Kiel, Federal Republik of Germany: 144-157.
- Rowland S. J., Allan G. L., Mifsud C., Nixon M., Boyd P., Glendenning D., 2005 Development of a feeding strategy for silver perch, *Bidyanus bidyanus* (Mitchell), based on restricted rations. *Aquac Res* 36: 1429-1441.
- Russell N. R., Fish J. D., Wootton R. J., 1996 Feeding and growth of juvenile sea bass: the effect of ration and temperature on growth rate and efficiency. *J Fish Biol* 49: 206-220.
- Silva C. R., Gomes L. C., Brandão F. R., 2007 Effect of feeding rate and frequency on tambaqui (*Colossoma macropomum*) growth, production and feeding costs during the first growth phase in cages. *Aquaculture* 264: 135-139.
- Sun L., Chen H., Huang L., Wang Z., 2006 Growth, faecal production, nitrogenous excretion and energy budget of juvenile cobia (*Rachycentron canadum*) relative to feed type and ration level. *Aquaculture* 259: 211-221.
- Tsevis N., Klaoudatos S., Conides A., 1992 Food conversion budget in sea bass, *Dicentrarchus labrax*, fingerlings under two different feeding frequency pattern. *Aquaculture* 101: 293-304.
- Yuan Y. C., Yang H. J., Gong S. Y., Luo Z., Yuan H. W., Chen X. K., 2010 Effects of feeding levels on growth performance, feed utilization, body composition and apparent digestibility coefficients of nutrients for juvenile Chinese sucker, *Myxocyprinus asiaticus*. *Aquac Res* 41: 1030-1042.
- Webster C. D., Thompson K. R., Muzinic L., 2002 Feeding fish and how feeding frequency affects sunshine bass. *J World Aquac Soc* 33: 20-24.
- Xiao-Jun X., Ruyung S., 1992 The bioenergetics of the southern catfish (*Silurus meridionalis* Chen): growth rate as a function of ration level and temperature. *J Fish Biol* 40: 719-740.

Received: 14 April 2012. Accepted: 16 June 2012. Published online: 22 June 2012.

Authors:

Majid Mohammad Nejad Shamoushaki, Department of Fishery, Bandar Gaz Branch, Islamic Azad University, Bandar Gaz, Iran, e-mail: majid_m_sh@bandargaziau.ac.ir

Zia Khari, Department of Fishery, Bandar Gaz Branch, Islamic Azad University, Bandar Gaz, Iran.

Zeynab Eslami, Department of Fishery, Bandar Gaz Branch, Islamic Azad University, Bandar Gaz, Iran.

How to cite this article:

Mohammad Nejad Shamoushaki M., Khari Z., Eslami Z., 2012 Determination of optimum feeding rate for growth of Caspian carp, *Cyprinus carpio* (Linnaeus, 1758) fingerlings. *AAFL Bioflux* 5(3):136-141.