

Influence of the protein quantity from combined fodder on the corporal development of guppy (*Poecilia reticulata*)

Adrian Gruber, Ioan M. Pop, Benone Păsărin, Gabriel Hoha, Cristian Radu-Rusu, Răzvan Radu-Rusu

University of Agricultural Sciences and Veterinary Medicine, Iasi, Romania. Corresponding author: A. Gruber, adygruber@yahoo.com

Abstract. This paper aims to highlight the effects of various concentrations of protein premixtures on the growth in length and weight of adult specimens of ornamental fish *Poecilia reticulata*, bred in captivity. Of this consideration, the experiment was set up with six groups: R1, R2, R3, TG, TP and TD, which were fed differently as follows: R1 with mixed fodder containing 30% CP (crude protein), own recipe; R2 with mixed fodder containing 30% GP, own recipe; R3 with mixed fodder containing 30% CP, own recipe; TG with Troco Grower fodder containing 43% CP; TP with Troco Prim fodder containing 44% CP; TD with Tetra Discus fodder containing 48% CP. The research was conducted over a period of three months in which there have been followed the length and weight, to determine body indices (average increase of the weight gain, Fulton index and average weight ratio multiplier).

Key Words: Guppy, *Poecilia reticulata*, mixed fodder, crude protein, body development, body indices.

Rezumat. Lucrarea are drept scop punerea în evidență a efectelor diferitelor concentrații de proteină din nutrețurile combinate administrate asupra creșterii în lungime și greutate a exemplarelor adulte din specia de pești ornamentali *Poecilia reticulata* crescute în captivitate. În acest sens, s-au format șase loturi experimentale: R1, R2, R3, TG, TP și TD care au fost hrănite diferențiat după cum urmează: R1 cu nutreț combinat după rețetă proprie cu 30% PB; R2 cu nutreț combinat după rețetă proprie cu 30% PB; R3 cu nutreț combinat după rețetă proprie cu 30% PB; TG cu nutreț Troco Grower cu 43% PB; TP cu nutreț Troco Prim cu 44% PB; TD cu nutreț Tetra Discus cu 48% PB. Cercetările s-au derulat pe o perioadă de trei luni în care s-au urmărit: lungimea și greutatea corporală pe baza cărora s-au determinat indicii corporali (șorul mediu de creștere în greutate, indicele Fulton și raportul de multiplicare a greutateii medii).

Cuvinte cheie: Guppy, *Poecilia reticulata*, nutreț combinat, proteină brută, dezvoltare corporală, indici corporali.

Introduction. Guppy (*Poecilia reticulata*) is one of the best model organisms used in various studies of sex-determination (Lindholm & Breden 2002; Lozinsky 2011; Mousavi-Sabet et al 2012), classical and molecular genetics (Dreyer et al 2007), color patterns (Petrescu-Mag & Bourne 2008), evolution of the color patterns (Hughes et al 1999), speciation and evolution (Alexander & Breden 2004), behavioral ecology (Houde 1997), spinal deformities (Gorman & Breden 2007), anatomical, physiological, embryological (Martyn et al 2006), ecotoxicological (Georgescu & Georgescu 2012; Talu et al 2012) and nutrition.

P. reticulata is a common exotic species raised in captivity conditions because it's high prolificacy (Bud 2006), low biological cycle (reaches sexual maturity at three months) and does not require special conditions for growth (Oprea 2000).

If when we talk about the consumption fish, it is mainly followed the relationship between body mass and skeletal weight, for aquarium fish the appearance and the body shape presents more interest. All this is determined by the quantity and quality of the given mixed fodder (Petrescu-Mag 2007).

This paper aims to highlight how the amount of protein in the fodder influences the body development of *P. reticulata*, one of the most common ornamental fish in

aquariums throughout the world. Proteins are substances of particular importance, they are only found in the highest proportion in organs and muscle tissue, performing biological functions that can not be met by another group of nutrients (Halga et al 2005).

Material and Method. Biological material on which the investigations were carried out was formed of 90 adults, aged 87-93 days, of the *P. reticulata* species, that were divided into six batches of 15 samples each. Each group was fed differently, as follows:

- group R1 with mixed fodder with 30% CP, own recipe;
- group R2 with mixed fodder with 30% CP, own recipe;
- group R3 with mixed fodder with 30% CP, own recipe;
- group TP with Troco Prim fodder with 43% CP;
- group TG with Troco Grower fodder with 44% CP;
- group TD with Tetra Discus fodder with 48% CP.

The research was conducted in the Laboratory of Aquaculture of USAMV Iasi, Romania, on a period of three months, time in which was followed the body development indices (increase in body weight and length).

For this purpose we used two tanks of the same capacity (15 liters) divided into three compartments each. Both tanks were fitted with the lighting (fluorescent lamp), air filter (electric indoor filter Hailea), portable aeration (aeration pump equipped with a nozzle Hailea), plant substrate (plants of the genus *Vallisneria* and *Elodea*), mineral substrate (quartz gravel). The two tanks were introduced in two aquariums, larger both in width and in length. The heating was performed with one electric 50W Kandege heater with thermostat, mounted outside the aquarium. The temperature reading was made using alcohol thermometers, one for each tank.

The followed parameters were average body length and average weight with which were calculated:

- Fulton index (maintenance index):

It was determined with the formula (Pășărin & Stan 2002):

$$L_i = \frac{g \times 100}{l^3}$$

where:

L_i = maintenance index;

g = weight (g);

l = standard body length (cm).

- Average increase weight gain:

It has been determined using the formula (Grether et al 2004):

$$S_p = G_f - G_i$$

where:

S_p = average gain of weight (mg);

G_f = average final weight (mg);

G_i = average initial weight (mg).

- The multiplying average weight ratio:

It was calculated with the formula (Misăilă et al 1990):

$$R.M.G.M. = \frac{G_f}{G_i}$$

where:

R.M.G.M. = the average weight multiplier;

G_f = average final weight;

G_i = average initial weight.

The parameters were determined at the end of each month throughout experiment. Body length was measured using an electronic calliper and weight using the Denver analytical balance type.

Results and Discussion. The gross chemical composition of the mixed fodder was given importance in the development of the fish's body. Each specie has specific nutrient requirements that must be provided by the administered combined fodder.

Table 1

The gross chemical composition of the used feeds

Combined fodder	DM (%)	A (%)	CP (%)	CF (%)	CC (%)
R1	93	13	30	7,0	4,4
R2	93	13	30	7,5	3,4
R3	93	13	30	8,0	4,0
Troco Prim	95	5	43	15,0	2,5
Troco Grower	93	10	44	22,0	1,1
Tetra Discus	94	11	48	6,5	2,0

DM – dry matter, A – ash, CP – crude protein, CF – crude fat, CC - crude cellulose.

The combined premixtures R1, R2 and R3 have the same amount of protein, respectively 30%, being obtained in the Quality Control Feed Laboratory of USAMV Iasi as were the nutritional requirements of the species *P. reticulata*. Commercial feeds have a higher percentage of protein which varied between 43-48%.




Figure 1. Evolution of average weight of the experimental groups.

For the specimen lots R1, R2 and R3, which were fed with compound feed with the same amount of protein (30%), the average weight had values close for each of the three months of experiment. Copies of lots TP, TG and TD achieved a higher growth in weight compared to previous batches due to higher amount of protein administered in the fodder (43%, 44% respectively 48%).

Table 2

The variation of the average body weight

Group	Specification	Month		
		1	2	3
R1	Variation interval (mg)	200 – 450	250 – 504	290 – 537
	\bar{x}	274.20	328.50	370.80
	$\bar{x} \pm s_{\bar{x}}$	274.20 ± 19.43	328.50 ± 19.22	370.80 ± 18.70
	C _v %	27.45	22.66	19.53
R2	Variation interval (mg)	120 – 420	178 – 478	228 – 525
	\bar{x}	210.80	269.00	318.50
	$\bar{x} \pm s_{\bar{x}}$	210.80 ± 19.34	269.00 ± 19.41	318.50 ± 19.16
	C _v %	35.53	27.94	23.30
R3	Variation interval (mg)	190 – 440	140 – 470	165 – 500
	\bar{x}	253.10	280.90	310.40
	$\bar{x} \pm s_{\bar{x}}$	253.10 ± 17.68	280.90 ± 20.32	310.40 ± 20.34
	C _v %	27.06	28.02	25.39
TG	Variation interval (mg)	210 – 980	310 – 1070	370 – 1110
	\bar{x}	484.30	588.60	649.10
	$\bar{x} \pm s_{\bar{x}}$	484.30 ± 64.14	588.60 ± 65.70	649.10 ± 64.72
	C _v %	51.29	43.23	38.62
TP	Variation interval (mg)	220 – 680	350 – 795	510 – 900
	\bar{x}	405	547.90	647.50
	$\bar{x} \pm s_{\bar{x}}$	405.00 ± 42.80	457.90 ± 37.47	647.50 ± 38.25
	C _v %	40.93	26.49	22.88
TD	Variation interval (mg)	220 – 610	340 – 725	445 – 830
	\bar{x}	395	513.10	621.90
	$\bar{x} \pm s_{\bar{x}}$	395.00 ± 32.88	513.10 ± 32.80	621.90 ± 32.79
	C _v %	32.23	24.76	20.42

The amount of protein directly influences the weight gain. The higher amount of protein in the feed, the greater is the increase of the weight (Figure 1, Table 2). However, the amount of the protein in the fodder should be rationally dosed, to avoid unjustified expenses with fodder, which would increase the final price and become economically onerous.

Statistics (Fisher and Tukey test) has been revealed that among the groups R1, R2 and R3 the differences in weight gain were insignificant throughout the experiment. Copies of these groups were fed with fodder obtained according to recipes R1, R2 or R3, containing the same amount of protein (30%). For lots TD, TP and TG the feed was with a higher amount of protein (43%, 44% and 48%), showing a greater increase in weight compared to groups R1, R2 and R3, these differences being distinctly significant. Between lots TD, TG and TP the statistical differences were insignificant.

Table 3

The evolution of the body weight in adults

Month	Statistical differences	Fisher / Tukey test	
1	SDD	$F_{\alpha 0,001} = 4.76 < F = 7.21$	
	DS	R1 – TD $w_{0,05} = 117.80 < w = 120.83 < w_{0,01} = 141.30$	
	ID	R1 – TP $w_{0,05} = 122.61 < w = 130.83 < w_{0,01} = 147.07$	
	SDD	R1 – TG $w = 210.12 > w_{0,01} = 141.30$	
	ID	R1 – R3 $w = 21.09 < w_{0,05} = 120.13$	
	ID	R1 – R2 $w = 63.40 < w_{0,05} = 120.13$	
	SDD	R2 – TD $w = 184.23 > w_{0,01} = 138.66$	
	SDD	R2 – TP $w = 194.23 > w_{0,01} = 144.10$	
	SDD	R2 – TG $w = 273.52 > w_{0,01} = 138.66$	
	ID	R2 – R3 $w = 42.31 < w_{0,05} = 117.80$	
	SDD	R3 – TD $w = 141.92 > w_{0,01} = 138.66$	
	SDD	R3 – TP $w = 151.92 > w_{0,01} = 144.10$	
	SDD	R3 – TG $w = 231.21 > w_{0,01} = 138.66$	
	ID	TG – TD $w = 89.29 < w_{0,05} = 113.51$	
	ID	TG – TP $w = 79.29 < w_{0,05} = 117.80$	
	ID	TP – TG $w = 10.00 < w_{0,05} = 117.80$	
	2	SDD	$F_{\alpha 0,001} = 4.14 < F = 13.2$
		SDD	R1 – TD $w = 184.57 > w_{0,01} = 135.81$
SDD		R1 – TP $w = 219.33 > w_{0,01} = 141.36$	
SDD		R1 – TG $w = 260.07 > w_{0,01} = 135.81$	
ID		R1 – R3 $w = 47.65 < w_{0,05} = 114.36$	
ID		R1 – R2 $w = 59.60 < w_{0,05} = 114.36$	
SDD		R2 – TD $w = 244.07 > w_{0,01} = 133.27$	
SDD		R2 – TP $w = 278.83 > w_{0,01} = 138.50$	
SDD		R2 – TG $w = 319.57 > w_{0,01} = 133.27$	
ID		R2 – R3 $w = 11.85 < w_{0,05} = 112.14$	
SDD		R3 – TD $w = 232.23 > w_{0,01} = 133.27$	
SDD		R3 – TP $w = 266.99 > w_{0,01} = 138.50$	
SDD		R3 – TG $w = 307.73 > w_{0,01} = 133.27$	
ID		TG – TD $w = 75.50 < w_{0,05} = 108.06$	
ID		TG – TP $w = 40.74 < w_{0,05} = 112.14$	
ID	TP – TG $w = 34.76 < w_{0,05} = 112.14$		
3	SDD	$F_{\alpha 0,001} = 3.33 < F = 27.62$	
	SDD	R1 – TD $w = 251.1 > w_{0,01} = 117.22$	
	SDD	R1 – TP $w = 276.75 > w_{0,01} = 119.64$	
	SDD	R1 – TG $w = 278.32 > w_{0,01} = 114.94$	
	ID	R1 – R3 $w = 60.37 < w_{0,05} = 96.79$	
	ID	R1 – R2 $w = 52.29 < w_{0,05} = 96.79$	
	SDD	R2 – TD $w = 303.4 > w_{0,01} = 114.94$	
	SDD	R2 – TP $w = 329.04 > w_{0,01} = 117.22$	
	SDD	R2 – TG $w = 330.61 > w_{0,01} = 112.79$	
	ID	R2 – R3 $w = 8.08 < w_{0,05} = 94.91$	
	SDD	R3 – TD $w = 311.47 > w_{0,01} = 114.94$	
	SDD	R3 – TP $w = 337.12 > w_{0,01} = 117.22$	
	SDD	R3 – TG $w = 338.69 > w_{0,01} = 112.79$	
	ID	TG – TD $w = 27.21 < w_{0,05} = 93.14$	
	ID	TG – TP $w = 1.57 < w_{0,05} = 94.91$	
ID	TP – TG $w = 25.64 < w_{0,05} = 96.79$		

SDD - significant distinct differences, DS - distinct differences, ID - insignificant differences.

Table 4

The evolution of body length in adults

Group	Specification	Month		
		1	2	3
R1	Variation interval (cm)	3.00 – 3.70	4.00 – 4.70	5.10 – 5.80
	\bar{x}	3.20	4.20	5.30
	$\bar{x} \pm s_{\bar{x}}$	3.20 ± 0.06	4.2 ± 0.05	5.3 ± 0.05
	$C_V\%$	6.64	4.87	3.87
R2	Variation interval (cm)	2.90 – 3.60	3.90 – 4.50	5.10 – 5.70
	\bar{x}	3.10	4.10	5.30
	$\bar{x} \pm s_{\bar{x}}$	3.10 ± 0.05	4.10 ± 0.05	5.30 ± 0.05
	$C_V\%$	5.75	4.36	3.38
R3	Variation interval (cm)	3.00 – 3.40	3.90 – 4.30	5.00 – 5.40
	\bar{x}	3.10	4.10	5.20
	$\bar{x} \pm s_{\bar{x}}$	3.10 ± 0.03	4.10 ± 0.03	5.20 ± 0.03
	$C_V\%$	4.17	3.15	2.48
TG	Variation interval (cm)	3.00 – 4.30	4.00 – 5.30	5.00 – 6.30
	\bar{x}	3.50	4.50	5.50
	$\bar{x} \pm s_{\bar{x}}$	3.50 ± 0.11	4.50 ± 0.11	5.50 ± 0.11
	$C_V\%$	12.57	9.50	7.79
TP	Variation interval (cm)	3.10 – 3.80	4.10 – 4.80	5.10 – 5.80
	\bar{x}	3.40	4.40	5.40
	$\bar{x} \pm s_{\bar{x}}$	3.40 ± 0.07	4.40 ± 0.07	5.40 ± 0.07
	$C_V\%$	8.19	5.89	4.80
TD	Variation interval (cm)	3.00 – 3.80	4.00 – 4.80	5.00 – 5.80
	\bar{x}	3.40	4.40	5.40
	$\bar{x} \pm s_{\bar{x}}$	3.40 ± 0.06	4.40 ± 0.06	5.40 ± 0.06
	$C_V\%$	7.31	5.60	4.57

The increase in length it is less influenced by the amount of protein administered in the combined fodder compared to the weight gain. Average body length had similar values in each of the three months of experimentation for all lots. In the first month there had been a body length ranged from 3.10 to 4.10 cm, in the second between 4.10 cm and 5.10 cm and in the third between 5.20 cm and 5.50 cm.




Figure 2. The evolution of the average body length.

From the previous chart we can see that the groups R1, R2 and R3 had similar values in the increase in length, the difference between them being 0.10 cm, while between lots TG, TP and TD the differences were larger (1 cm).

In the second month the increase in length was 1 cm for all the experimental groups compared to the first month, while in the third month the growth varied between 1.10 cm and 1.30 cm, slightly higher values for the groups TP, TG and TD.

The amount of protein in the given fodder is one of the factors determining the growth in length, with environmental conditions (temperature, pH, water chemistry) and social factors (stocking density, relations between samples in the group).

At the end of the three months of experimentation there was a varied value of the average length in the lots fed with the laboratory recipes from 5.20 cm for group R3 and to 5.30 cm for R1 group, while the groups fed with commercial feed had body length values between 5.40 cm and 5.60 cm.

The average body length for lots R1, R2 and R3 had similar values at the end of each experiment; the differences between groups were not significant, a possible explanation could be that the feeds used had the same concentration of protein, of 30%. For the groups fed with Troco Prim and Tetra Discus, the average body length value at the end of three months of experimentation was 5.40 cm, although they received fodder with different concentrations of protein (43% for Troco Prim and 48% for Tetra Discus). The average body length for the TG group, whose copies were fed with Troco Grower (44% CP) was with 0.20 cm higher than the TD group, which was fed with Tetra Discus (48% CP).

From this we conclude that the value of the body length is influenced in a lesser extent of the amount of protein in the administered fodder, compared to the weight gain.

Table 5

The evolution of body length in adults

Month	Statistical differences	Fisher / Tukey test
1	SDD	$F_{\alpha 0,01} = 3.34 < F = 3.95 < F_{\alpha 0,001} = 4.76$
	ID	$R1 - TD w = 0.14 < w_{0,05} = 0.22$
	ID	$R1 - TP w = 0.15 < w_{0,05} = 0.23$
	SD	$R1 - TG w_{0,05} = 0.22 < w = 0.24 < w_{0,01} = 0.26$
	ID	$R1 - R3 w = 0.12 < w_{0,05} = 0.22$
	ID	$R1 - R2 w = 0.10 < w_{0,05} = 0.22$
	SD	$R2 - TD w_{0,05} = 0.22 < w = 0.25 < w_{0,01} = 0.26$
	SD	$R2 - TP w_{0,05} = 0.22 < w = 0.25 < w_{0,01} = 0.27$
	SDD	$R2 - TG w = 0.34 > w_{0,01} = 0.26$
	ID	$R2 - R3 w = 0.02 < w_{0,05} = 0.22$
	SD	$R3 - TD w_{0,05} = 0.22 < w = 0.26 < w_{0,01} = 0.27$
	SD	$R3 - TP w_{0,05} = 0.22 < w = 0.27 < w_{0,01} = 0.28$
	SDD	$R3 - TG w = 0.36 > w_{0,01} = 0.26$
	ID	$TG - TD w = 0.09 < w_{0,05} = 0.21$
	ID	$TG - TP w = 0.09 < w_{0,05} = 0.22$
	ID	$TP - TD w = 0.01 < w_{0,05} = 0.22$
	2	SDD
ID		$R1 - TD w = 0.15 < w_{0,05} = 0.21$
ID		$R1 - TP w = 0.16 < w_{0,05} = 0.21$
SDD		$R1 - TG w = 0.29 > w_{0,01} = 0.25$
ID		$R1 - R3 w = 0.14 < w_{0,05} = 0.21$
ID		$R1 - R2 w = 0.10 < w_{0,05} = 0.21$
SDD		$R2 - TD w = 0.25 > w_{0,01} = 0.24$
SDD		$R2 - TP w = 0.26 < w_{0,01} = 0.25$
SDD		$R2 - TG w = 0.39 > w_{0,01} = 0.24$
ID		$R2 - R3 w = 0.04 < w_{0,05} = 0.21$
SDD		$R3 - TD w = 0.29 > w_{0,01} = 0.24$
SDD		$R3 - TP w = 0.30 > w_{0,01} = 0.25$
SDD		$R3 - TG w = 0.43 > w_{0,01} = 0.24$
ID		$TG - TD w = 0.14 < w_{0,05} = 0.20$
ID	$TG - TP w = 0.13 < w_{0,05} = 0.21$	
ID	$TP - TD w = 0.01 < w_{0,05} = 0.21$	
3	ID	$F = 1.5 < F_{\alpha 0,05} = 2.37$

SDD - significant distinct differences, SD - significant differences, ID - insignificant differences.

In the first month, the maintenance index had values around 1, the specimens having a good maintenance. In the next two months, the relationship between body weight and length was reduced because some females entered in the gestation period and therefore had a greater weight gain compared to the increase in length. Weight gain of the females is illustrated also by the increase of weight gain, being bigger for the lots TG, TP and TD.

In the second month the maintenance index decreased for all the experimental groups, being minimum for lot R2 (0.31) and maximum for TP and TD groups (0.74). The length and weight growth decreased with age. Therefore, also the Fulton index's values were lower than the first month.

The multiplication average weight ratio had similar values for all groups, ranging between 1.11 and 1.35.

In the third month, the maintenance index declined further for all the experimental groups, being minimum for lot R2 (0.21) and maximum for the TP group (0.41). Weight gain during the gestation period at females resulted in a greater weight gain for lots TG, TP and TD, compared to the groups R1, R2 and R3. The multiplication of the average weight ratio for all groups had similar values, ranging between 1.11 and 1.21.

Table 6

Corporeal indices in adults

Specification	Month	Experimental group					
		R1	R2	R3	TP	TG	TD
Average body length (cm)	1	3.20	3.10	3.10	3.20	4.10	3.10
	2	4.20	4.10	4.10	4.20	5.10	4.10
	3	5.30	5.30	5.20	5.40	5.50	5.40
Average initial weight (mg)	1	216.0	157.8	228.5	278.8	333.3	273.3
	2	274.2	210.8	253.1	405.0	484.3	395.0
	3	328.5	269.0	280.9	547.9	588.6	513.1
Average final weight (mg)	1	274.2	210.8	253.1	405.0	484.3	395.0
	2	328.5	269.0	280.9	547.9	588.6	513.1
	3	370.8	318.5	310.4	647.5	649.1	621.9
Maintenance index (Fulton)	1	0.84	0.71	0.85	1.03	1.13	1.01
	2	0.44	0.31	0.41	0.74	0.44	0.74
	3	0.25	0.21	0.22	0.41	0.39	0.40
Average weight gain (mg)	1	58.2	53.0	24.6	126.2	151.0	125.2
	2	54.3	58.2	27.8	142.9	104.3	118.1
	3	42.3	49.5	29.5	99.6	60.5	118.8
Multiplication ratio of average weight	1	1.27	1.34	0.90	1.45	1.45	1.45
	2	1.20	1.28	1.11	1.35	1.22	1.30
	3	1.13	1.18	1.11	1.18	1.10	1.21

Conclusions

1. The weight gain is influenced directly proportional by the percentage of protein in the given combined fodder;
2. The percentage of protein in combined fodder used for feeding the fish of the species *P. reticulata* causes small changes in body length;
3. The maintenance index's values increase proportionally to the percentage of the protein in the used combined fodder;
4. Average value of the weight gain is directly dependent on the protein amount of the used food;
5. Body development of the species of *P. reticulata* depends on the gross content in protein of the used mixed fodder.

Acknowledgements. This work was co financed from the European Social Fund through Sectorial Operation Programme, Human Resources Development 2007-2013, project number POSDRU/1.89/1.5/S62371 Postdoctoral School in Agriculture and Veterinary Medicine.

References

- Alexander H. J., Breden F., 2004 Sexual isolation and extreme morphological divergence in the Cumaná guppy: a possible case of incipient speciation. *J Evol Biol* 17:1238-1254.
- Bud I., 2006 [Small aquarium encyclopedia]. Risoprint Publishing House, Cluj-Napoca. [In Romanian].
- Dreyer C., Hoffmann M., Lanz C., Willing E. M., Riester M., Warthmann N., Sprecher A., Tripathi N., Henz S., Weigel D., 2007 ESTs and EST-linked polymorphisms for genetic mapping and phylogenetic reconstruction in the guppy, *Poecilia reticulata*. *BMC Genomics* 8:269.
- Georgescu B., Georgescu C. E., 2012 *Poecilia reticulata* as a valuable biological indicator of endocrine disruption. *Poec Res* 2(1):15-19.
- Gorman K. F., Breden F., 2007 Teleosts as models for human vertebral stability and deformity. *Comp Biochem Physiol C Toxicol Pharmacol* 145(1):28-38.

- Grether G. F., Kasahara S., Kolluru G.R., Cooper E.L., 2004 Sex-specific effects of carotenoid intake on the immunological response to allografts in guppies (*Poecilia reticulata*). Proc R Soc Lond B 271:45-49.
- Halga P., Pop I. M., Avarvarei T., Popa V., 2005 [Animal nutrition and feeding]. Alfa Publishing House, Iasi. [In Romanian].
- Hughes K. A., Du L., Rodd F. H., Reznick D. N., 1999 Familiarity leads to female mate preference for novel males in the guppy, *Poecilia reticulata*. Anim Behav 58:907-916.
- Hoode A. E., 1997 Sex, color, and mate choice in guppies. Princeton University Press, New Jersey.
- Lindholm A., Breden F., 2002 Sex chromosome and sexual selection in Poeciliid fishes. Am Nat 160:S214-S224.
- Lozinsky L. R., 2011 [New contributions to sex and body size control in guppy fish (*Poecilia reticulata*)]. Poec Res 1(1):1-19. [In Romanian].
- Martyn U., Weigel D., Dreyer C., 2006 In vitro culture of embryos of the guppy, *Poecilia reticulata*. Dev Dyn 235:617-622.
- Misăilă C., Misăilă R. E. et al 1990 [Incorporation of the tuff and procaine in the rainbow trout (*Salmo gairdneri* Rich.) fodder]. "Piscicultura Moldovei" Scientific Papers 1:223-230. [In Romanian].
- Mousavi-Sabet H., Langroudi H. F., RohaniRad M., 2012 Sex reversal, mortality rate and growth of guppy (*Poecilia reticulata*) affected by 17-alpha methyltestosterone. Poec Res 2(1):1-8.
- Oprea L., 2000 [Fish nutrition and feeding]. Orizonturi Universitare Publishing House, București. [In Romanian].
- Pășărin B., Stan T., 2002 [Aquaculture – practical guide]. Pim Publishing House, Iași. [In Romanian].
- Petrescu-Mag I. V. (ed), 2007 [Sex control in guppyculture]. Academicpres, Cluj-Napoca [in Romanian].
- Petrescu-Mag I. V., Bourne G. R., 2008 Crossing-over between Y chromosomes: another possible source of phenotypic variability in the guppy, *Poecilia reticulata* Peters. AACL Bioflux 1(1):1-10.
- Talu S., Petrescu-Mag I. V., Pasarin B., 2012 Investigation on acute toxicity of lindane in guppies, *Poecilia reticulata* Peters, 1859. Poec Res 2(1):9-14.

Received: 18 October 2012. Accepted: 20 November 2012. Published online: 12 February 2013.

Authors:

Adrian Gruber, University of Agricultural Sciences and Veterinary Medicine, Faculty of Animal Science, Romania, Iasi, M.Sadoveanu Alley, no.8, adygruber@yahoo.com;

Ioan Mircea Pop, University of Agricultural Sciences and Veterinary Medicine, Faculty of Animal Science, Romania, Iasi, M.Sadoveanu Alley, no.8, popim@univagro-iasi.ro;

Benone Pășărin, University of Agricultural Sciences and Veterinary Medicine, Faculty of Animal Science, Romania, Iasi, M.Sadoveanu Alley, no.8, pbeno@univagro-iasi.ro;

Gabriel Hoha, University of Agricultural Sciences and Veterinary Medicine, Faculty of Animal Science, Romania, Iasi, M.Sadoveanu Alley, no.8, gabihoha@yahoo.com;

Cristina Radu-Rusu, University of Agricultural Sciences and Veterinary Medicine, Faculty of Animal Science, Romania, Iasi, M.Sadoveanu Alley, no.8, mimicris@yahoo.com;

Răzvan Radu-Rusu, University of Agricultural Sciences and Veterinary Medicine, Faculty of Animal Science, Romania, Iasi, M.Sadoveanu Alley, no.8, rprobios@yahoo.com.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Gruber A., Pop I. M., Benone P., Hoha G., Radu-Rusu C., Radu-Rusu R., 2013 Influence of the protein quantity from combined fodder on the corporal development of guppy (*Poecilia reticulata*). AACL Bioflux 6(2):145-154.