



Authentication of *Tenualosa* species in Perak River, Malaysia: application of morphological measurement and molecular analysis of partial *CO1* and *16S* genes to resolve species ambiguity

¹Nurul N. Arjunaidi, ²Muhammad F. Zakaria, ¹Abdul H. Abdul Aziz, ^{1,3}M. Sheriff Shahreza, ^{1,3}Tun N. Aimi Mat Jaafar, ^{1,4}Y. Giat Seah, ^{1,3}Ariffin Nur Asma

¹ School of Fisheries and Aquaculture Sciences, Universiti Malaysia Terengganu, Terengganu, Malaysia; ² School of Marine Science and Environment, Universiti Malaysia Terengganu, Terengganu, Malaysia; ³ Institute of Tropical Aquaculture, Universiti Malaysia Terengganu, Terengganu, Malaysia; ⁴ Fish Division South China Sea Repository and Reference Center, Institute of Oceanography and Environment, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia.
Corresponding author: A. Nur Asma, nurasma@umt.edu.my

Abstract. Malaysia possessed two species of *Tenualosa* sp.; the *Tenualosa tolli* and *Tenualosa macrura*, which can only be found in Sarawak. Recently, samples that resembled the *Tenualosa* sp. were found in Perak River of Peninsular Malaysia. The absence of comprehensive study of *Tenualosa* sp. in Peninsular Malaysia had urged this study to be conducted to authenticate the samples found. Identification were done using morphological analysis with the integration of molecular assessment using partial genes of cytochrome c oxidase subunit 1 (*CO1*) and 16S rRNA. Morphological analysis of 40 samples showed the closest correlation with *Tenualosa ilisha* (Hamilton, 1822). However, variations were found in meristic counts and the samples were differentiated into two groups. The meristic counts also overlapped with other species of the same genus. Molecular assessments on 10 samples performed using partial *CO1* and 16S rRNA genes had obtained 500-bp and 573-bp sequences respectively, that has high similarity with *T. ilisha*. Further genetic assessment through phylogenetic analysis showed that all 10 samples formed a monophyletic lineage with *T. ilisha*. However, the samples were significantly separated with others of the same species from different countries. The findings suggested that there is a possibility the samples might migrate from other countries and had experienced domestic speciation towards the local environment. The findings of this study would be useful in recognizing the potential fisheries resources. This will greatly help in planning and implementing a healthy and sustainable exploitation of this species.
Key Words: *Tenualosa* sp., morphological assessments, cytochrome C oxidase subunit 1 gene, 16S rRNA gene.

Introduction. *Tenualosa* species or known as Terubok is an important and high commercial value clupeid fish. There are five species of *Tenualosa* spp. can be found worldwide and two species are common in Malaysia. These two species are *Tenualosa tolli* and *Tenualosa macrura* which can be found only in Sarawak waters (Blaber et al 1996; Phillip 2001). The Sarawak *Tenualosa* species is one of the most significant estuarine fish that is considered as one of the Malaysian commercial fish species that contributes to the national economy. In Peninsular Malaysia, catching of *Tenualosa* species is rare and inconsistent in the past few decades (Gambang 1988). It has been presumed that the *Tenualosa* species has subsequently extinct, however no such records has ever been confirmed. Recently, catch landings of *Tenualosa* species in Peninsular Malaysia has been proclaimed by several local fishermen in Perak River. The reappearance of *Tenualosa* species in Perak River has known to be disappeared for almost 20 years has raised several research questions such as its distribution, taxonomy, and systematics. However, *Tenualosa* species in Perak River has never received enough attention, hence

no comprehensive studies are done. Consequently, there is lack of information regarding the *Tenualosa* species in Perak River. To date, this species is still unidentified and there is no current study to officially validate its species even though the *Tenualosa* species in Perak bare resemblance with the *Tenualosa* species in Sarawak. Thus, the current status of *Tenualosa* species in Perak River is still poorly understood.

As the identification of this species that are caught in Perak River remains unclear, this might lead to the difficulty of collecting any information regarding the species. Insufficient knowledge and information may leave the *Tenualosa* species in Perak unattended without any efforts of its preservation and conservation management in Perak River. Pollution and overfishing might occur and could lead to unsustainable exploitation and the declination in numbers of this species that might render its extinction. Such incidents have been observed in the other two species; the *T. toli* and *T. macrura* in Sarawak as the numbers have been reported drastically decreasing in the past few years (Phillip 2001; Rahim et al 2014). Abdul Aziz et al (2015) reported that genetic diversity of *T. toli* was relatively low for two populations in Sarawak and might cause a genetic deprivation towards its population. Therefore, as an initial step, identifying the species in Perak River will acknowledge the local community and authorities regarding the value of the species and efforts to manage the species can be implemented. Moreover, *Tenualosa* species in Perak River can become an alternative source of Malaysian *Tenualosa* species supply besides Sarawak. Herein, it is hypothesized that the *Tenualosa* species in Perak River is of different species from the *T. toli* and *T. macrura* from Sarawak.

One of the best ways to precisely identify species is through morphometric study accompanied by molecular identification. Kartavtsev et al (2006) reported that molecular approach is more accurate in species identification compared to morphological character identification alone. Several studies have been done through the application of both approaches to precisely validate and identify the aquatic species (Watanabe et al 2008; Kolangi-Miandare et al 2013; Haniffa et al 2014). The development of species-specific morphological and molecular markers has been done to successfully monitor natural population stocks (Tinti et al 2003). It has been demonstrated that both markers can be applied to investigate the species divergence and genetic variation within geographical regions, hence relatively mapping the distribution (Motamedi et al 2014). As a result, efforts to manage natural fisheries population stocks can be implemented and applied to maintain sustainability. Such studies have been done in natural populations of *Rastrelliger* spp. and *Channa striata* in Malaysia (Darlina et al 2011; Song et al 2013). However, as any information regarding the *Tenualosa* species in Perak, identification has been poorly documented, the efforts to manage the natural populations cannot be done. This might cause unsustainable exploitations of fisheries resources and once again causing the disappearance of valuable *Tenualosa* species resources in the future. Therefore this study was conducted to identify the *Tenualosa* species in Perak through morphological and molecular approaches.

Material and Method. Samples of *Tenualosa* species were collected from the Perak River in Bagan Datoh, Perak from January - March 2014. Samples were caught using drift net by the fishermen. A total of 40 samples were collected and used for the experiment. White muscle tissue and fin clips were collected and preserved in 95% ethanol and stored at -20°C for further genetic analysis. The tissue and clips were taken on the right side of the samples to allow the left side to be used for morphological analysis.

Morphological analysis. The samples were identified morphologically according to the verification of Narejo et al (2008), Munroe et al (1999) and Hong et al (2013). Twenty five morphometric characters were measured using digimatic digital caliper (Mitutoyo, Japan) with an accuracy of 0.01 mm and seven meristic characters were counted using standard methods. The procedure of fixation, photograph and preservation for the specimen are followed Seah et al (2011). Methods of counting and measuring are generally followed Hubbs & Lagler (1964) and terminology of morphological features and descriptions are modified from Seah et al (2009). Measurements of the morphometric

characters were standardized in order to eliminate any size effect: Standardized measurement = Morphometric Parameter / Standard Length*100.

DNA extraction, Polymerase Chain Reaction (PCR) amplification and sequencing. Total genomic DNA of 10 samples were extracted using DNA extraction kits, the Cell/Tissue DNA Extraction Kit (Spin Column) (BioTeke Corporation, China). Two molecular markers were used, the partial cytochrome c oxidase subunit 1 (*CO1*) and 16S gene marker.

The *CO1* gene was amplified using two primers based on Jannatul-Fariyah et al (2011) which were CO1f (5'-CCTGCAGGAGGAGGAGAYCC-3') and CO1r (5'-CCAGAGATTAGACCGAAATCAGTG-3'). A PCR mixture containing 2.5 µL of 10× PCR buffer; 0.8 µL of 25 mM MgCl₂; 0.6 µL of 100 pmol of both primers; 0.4 µL of 5U Taq DNA polymerase (Vivantis, Malaysia); 2.4 µL of 10,000 uM dNTP; and 3.0 µL of 100 ng DNA sample was prepared in a 25 µL reaction. The cycling profile for *CO1* began with the initial step of 3 minutes at 94°C, followed by 30 cycles of denaturation (at 94°C for 30 seconds), annealing (at 52°C for 40 seconds) and extension (at 72°C for 60 seconds) and finally subjected to the final extension (at 72°C for 10 minutes).

The 16S gene was amplified using primers 16SarL (5'- CGCCTGTTTATCAAAAACAT -3') and 16SbrH (5'- CCGGTCTGAACTCAGATCACG T-3') (Palumbi 1996). A PCR reaction with a total volume of 25 µL were prepared using 2.5 µL of 10× PCR buffer (readily added with MgCl₂); 0.3 µL of 10 µM of both primers; 0.2 µL of 5U Taq DNA polymerase (TransGen Biotech Co., Ltd, China); 0.25 µL of 10,000 uM dNTP; and 4.0 µL of 100 ng DNA sample. PCR cycling conditions used were as follow: initial denaturation at 94°C for 2 min, 30 cycles of denaturation at 93°C for 30 s, primer annealing at 55°C for 40 s and chain elongation at 72°C for 60 s, follow by final extension at 72°C for 5 min. Both the amplifications were done in a Thermal Cycler (Eppendorf AG, Germany).

Prior to sequencing, the PCR products were purified using a DNA purification kit (BioTeke Corporation, China) according to the manufacturer instruction. All the purified amplicons were then sequenced bi-directionally by First Base Laboratories Sdn. Bhd. using the same primers for PCR amplification.

Sequence alignment and evolutionary analysis. A Macintosh-based software; eBioX (<http://www.ebioinformatics.org/ebiox/>) were used to edit the obtained sequences. The chromatograms were analyzed and inspected for nucleotides conformity using the software 4Peaks (Nucleobytes Inc.) and FinchTV (Geospiza Inc.). The processes were to remove any unwanted sequences, noise, gaps and stop codons to precisely validate the sequences. The sequences were then identified using Nucleotide Basic Local Alignment Search Tool (BLAST) (Altschul et al 1990) at the National Center of Biotechnology Information (NCBI) database (<http://www.ncbi.nlm.nih.gov>). Next, the edited sequences were aligned together with sequences of other *Tenuulosa* species available in the GenBank (*T. toli*, Accession number: AP011600.1; *T. thibaudeaui*, Accession number: AP011604.1). *T. ilisha* from other countries such as from India, Thailand and West Africa were also included (*T. ilisha* Thailand, Accession number: AP011611.1; *T. ilisha* India 1, Accession number: AP011610.1; *T. ilisha* India 2, Accession number: NC016682.1; *T. ilisha* India 3, Accession number: DQ400344.1; *T. ilisha* West Africa, Accession number: EU552785.1). This is to assist in inferring the possibility of evolutionary origin of the samples found in Perak River.

The alignment was performed in MEGA software through CLUSTAL W program (version 5.2) (Tamura et al 2011). *Lutjanus argentimaculatus* (Accession number: DQ400344.1) and *Cromileptes altivelis* (Accession number: KC845547.1) were used as outgroup. Phylogenetic trees of Neighbour-Joining, Maximum-Parsimony and Maximum-Likelihood were constructed to resolve the evolutionary lineage amongst the *Tenuulosa* species. A best-fit DNA substitution model of Kimura-2-parameter was used in NJ and ML analyses with the confidence level of 1000 replications (Tamura et al 2011).

Results and Discussion. Morphometric and meristic analysis of *Tenuulosa* species collected from Perak River revealed that more than 90% of the samples had been identified as *T. ilisha*. Out of 25 morphometric and seven meristic characters, five morphometric and four meristic characters were chosen to identify the samples collected

significantly. Morphological assessments showed that the samples could be differentiated into two groups based on the variations of morphometric measurement and meristic count (Table 1). The average total length and standard length \pm S.D for group A were 30.6 ± 2.59 and 24.3 ± 1.78 respectively whereas for group B were 30.2 ± 1.71 and 22.8 ± 1.36 respectively. Narejo et al (2008) made a comparison between two types of *T. ilisha* by grouping and differentiated the morphometric and meristic characters based on size range. The result of Total Length (TL), Standard Length (SL) and Fork Length (FL) and meristic of the study had been used to compare with the results of present study to assist in identifying the *Tenualosa* species in Perak River.

Table 1
Comparison of *Tenualosa ilisha* morphological characters between the two groups (Group A and Group B) and *Tenualosa ilisha* from Pakistan (Type A and Type B)

Morphometric	<i>Tenualosa</i> sp. (Group A)		<i>Tenualosa</i> sp. (Group B)		<i>Tenualosa ilisha</i> (Narejo et al 2008)			
	Min-Max	Mean \pm S.D.	Min-Max	Mean \pm S. D.	Type A		Type B	
					Length groups	Length groups	Length groups	Length groups
					25.1-30.0*	30.1-35.0*	25.1-30.0*	30.1-35.0*
					Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.
Total length (TL) (cm)	27.4-34.8	30.6 \pm 2.59	27.4-34.1	30.2 \pm 1.71	28.0 \pm 1.90	32.1 \pm 2.10	27.0 \pm 1.55	33.4 \pm 1.55
Standard length (SL) (cm)	22.4-27.2	24.3 \pm 1.78	20.0-25.3	22.8 \pm 1.36	22.2 \pm 1.33	26.0 \pm 1.88	21.5 \pm 1.20	27.0 \pm 1.55
Fork length (FL) (cm)	25.0-30.0	26.9 \pm 1.97	20.8-28.6	25.6 \pm 2.02	23.5 \pm 1.88	27.5 \pm 1.33	23.0 \pm 1.22	28.5 \pm 1.15
Head length (HL) (cm)	6.4-7.8	6.9 \pm 0.43	5.0-7.4	6.5 \pm 0.49	6.1 \pm 0.80	7.3 \pm 0.65	5.3 \pm 0.60	5.5 \pm 0.33
(HL as % to SL)	26.6-31.0	28.6 \pm 1.09	20.2-32.4	28.7 \pm 1.93	-	-	-	-
Eye diameter (ED) (cm)	1.0-1.5	1.2 \pm 0.13	1.0-1.5	1.2 \pm 0.11	1.1 \pm 0.55	1.2 \pm 0.44	1.0 \pm 0.25	1.0 \pm 0.22
(ED as % to SL)	4.3-5.5	5.0 \pm 0.41	4.6-6.4	5.3 \pm 0.45	-	-	-	-
Meristic	Range	Mean \pm S.D.	Range	Mean \pm S.D.	Range	Mean \pm S.D.	Range	Mean \pm S.D.
Dorsal fin rays	18-20	-	16-18	17 \pm 1.0	17-19	18 \pm 1.0	18-20	19 \pm 1.0
Pectoral fin rays	14-15	-	12-15	13 \pm 1.0	15-16	14 \pm 1.0	14-15	13 \pm 1.0
Pelvic fin rays	8	-	7-9	8 \pm 0.0	7-9	8 \pm 1.0	8-10	9 \pm 1.0
Anal fin rays	19-21	-	15-20	18 \pm 1.0	18-22	20 \pm 2.0	19-23	21 \pm 1.0

*Length groups (cm) of samples used, taken from Narejo et al (2008).

Results of the present study showed that the HL as percentage of SL for group A and group B were 28.6% and 28.7%, respectively. Whitehead (1985) reported that the HL for *T. ilisha* is 28 to 32% of its SL, for *T. macrura* is 22 to 25% of SL, for *T. reevesii* is 27 to 33% of SL, for *T. thibaudeaui* is 30 to 33% of SL and *T. toli* is 25 to 27% of SL. However, as results showed are overlapping to more than one species of *Tenualosa*, the *Tenualosa* species from Perak cannot be accurately identified. Nevertheless, the range of meristic characters of both groups was quite similar to Narejo et al (2008). Narejo et al (2008) found that the *T. ilisha* in Pakistan have 17-20 dorsal fin rays (DFR), 14-16 pectoral fin rays (PCFR), 7-10 pelvic fin rays (PVFR), 18-23 anal fin rays (AFR). As in this study, the Group A showed the DFR was 18-20, PCFR was 14-15, PVFR was 8 and the AFR was 19-21. However, *Tenualosa* species from Perak - group B has meristic characters that are slightly out of range and not in consensus with Narejo et al (2008). Group B showed the DFR was 17 ± 1.0 , PCFR was 13 ± 1.0 and AFR was 18 ± 1.0 , these are less than those from Group A and Pakistan *T. ilisha*. Arai & Amalina (2014) recorded the presence of *T. ilisha* in Malaysian waters by examining fourteen samples using morphological characters only. The results of the study reported that the *T. ilisha* has a measurement HL of 26.5-30.0%, which concurrent with the present study. However, for the meristic characters, Arai & Amalina (2014) reported that the DFR is 17-18, PCFR is 13-15, PVFR is 8, and AFR is 18-21, which is quite variable than the result of the present study. Forty samples are


examined in the present study and the morphological characters are ambiguous. Thus, the results indicate that there are morphological variations amongst the *Tenualosa* species in Perak. Variations of morphological characters of the specimens might be related to the point of origin and the aquatic ecosystems they thrived in. Study by Narejo et al (2008) emphasized that the length-weight relationship of *T. ilisha* was found to be significantly different in accordance to the winter-summer seasons.

Variations in morphological characters have also been observed in populations of *Puntius bimaculatus* in terms of physio-chemical parameters within different altitudinal range of river basin (De Silva & Liyanage 2009). Furthermore, morphological variations amongst marine fish species have been documented in Baltic Sea herring (*Clupea harengus*) (Jorgensen et al 2008) and bluefish (*Pomatomus saltatrix*) (Turan et al 2006). Both studies were done by examining samples throughout oceanic system with no physical barriers, thus suggesting the possibility of occurrence of migration. Turan et al (2006) suggested that variations in morphometric and meristic characters might also be influenced by several factors such as different feeding environment, prey types and food availability due to different environmental factors. Turan et al (2006) also exerted that differentiation in phenotypic traits do not tend to hinder gene flow between populations but also implied that the fish groups might not mix extensively. Present study indicates that there is a possibility that Group A of *Tenualosa* species in Perak might migrate from the Indian Ocean whereas Group B might be a later generation of *T. ilisha* of Group A. However, further studies should be done to confirm this matter.


The variations of morphological characters in *Tenualosa* species in Perak has led to the used of molecular tools to genetically identify the samples. Two molecular markers of *CO1* and 16S rRNA has been employed to determine the identity of *Tenualosa* species found in Perak River. Sequence analysis of the amplified fragments of partial *CO1* and 16S revealed a sequence of 500-base-pair and 573-base-pair, respectively for each samples. The *CO1* and 16S sequences were deposited in GenBank under accession numbers KP128692-KP128696 (*CO1*) and KP128687-KP128691 (16S). BLAST results showed that all samples had 99% similarity with mitochondrial DNA *CO1* and 16S sequences of *T. ilisha* (Accession Number: AP011611.1) in the GenBank Database, authenticating that all the samples are *T. ilisha*. The low genetic distance (*CO1*: < 0.017, 16S rRNA: < 0.028) between *Tenualosa* species in Perak and other *T. ilisha*, further support the species authentication. Phylogenetic tree of Neighbour-Joining (NJ), Minimum-Evolution (ME), Maximum-Parsimony (MP) and Maximum-Likelihood (ML) showed similar patterns in which all samples of *Tenualosa* species in Perak were separated from the sister group and outgroup and resolved within the same monophyletic clade with other *T. ilisha*.

Interestingly, all samples of *Tenualosa* species in Perak formed a single distinctive clade that is independent from other lineage of *T. ilisha* from Thailand, India and West Africa (Figure 1). Both genes separated the *Tenualosa* species in Perak with strong statistical support (100%). This suggested that there is a possibility the *Tenualosa* species in Perak is a distinctive and isolated species of *T. ilisha* that can be differentiated from other fishes of the same species. In addition, the evolutionary divergence of *Tenualosa* species in Perak from its monophyly lineage might also indicate that they also share a common ancestor. A comprehensive study should be done in the future involving a wider geographical range and higher number of samples. By doing so, investigation of genetic structuring might confirm the hypothesis. Therefore, the origin and evolutionary history of *Tenualosa* species in Perak could be possibly determined. Such case has been documented on the Telmatherinidae fishes (Stelbrink et al 2014). Study done on *Sphaeramia orbicularis* demonstrated the occurrence of genetic differentiation in populations of isolated fishes of the same species (Hanzawa et al 2012). Each population will have its own unique haplotypes. However, a common haplotype shared between several populations indicates the possibility of a closer inter-population relationship through a common ancestor in which has been documented in studies done on *Feneropenaeus chinensis* (Li et al 2009), *Channa striata* (Siti-Balkhis et al 2011), *Odontesthes* spp. (Garcia et al 2014) and *Tenualosa toli* (Abdul Aziz et al 2015). Plath et al (2010) in their study on Atlantic molly (*Poecilia mexicana*), suggested that speciation

might occur without the limitation of geographical barriers and through domestic adaptation to different environment.



(a)



(b)

Figure 1. Phylogenetic position of *Tenualoosa* sp. from Perak River of Peninsular Malaysia compared to Thailand, India and West Africa *CO1* (a) and 16S rRNA (b) gene sequences. Neighbour-Joining tree constructed using Neighbour-Joining algorithm with Kimura-2 parameter modal. Value at each node is the bootstrap value (1000 replicate). Genbank accession numbers are provided after the species name. *Cromileptes altivelis* (Serranidae) and *Lutjanus argentimaculatus* (Lutjanidae) are used as outgroup. Specimens in the presence study are labeled with the initial TP (*Tenualoosa* species in Perak).

Further study revealed that local adaptation towards varied environment greatly influences the molecular characteristics (Kelley et al 2012). *T. ilisha* has been reported to have great capability of long distance migration (up to 700-1200km) (Pillay & Rosa 1963; McDowall 1988). Hence, there is a possibility of migration and local adaptation of *T. ilisha* of the Indian Ocean through several different countries as there is limitless connectivity of the oceanic systems. The results of genetic assessment are congruent with the

morphological analysis. In contrary to study done by Arai & Amalina (2014), meristic counts investigated in forty samples of *Tenuulosa* species found in Perak River exhibits slight variation from those of *T. ilisha*. This might possibly correlate with the evolutionary lineage that showed significant separation of *Tenuulosa* species found in Perak River with the other *T. ilisha*. The findings of the present study would assist in acknowledging the existence of *T. ilisha* in Malaysia. This would help in designing and implementing conservation programs in managing these fisheries resources in the future.

Conclusions. The present study has successfully identified *Tenuulosa* species in Perak of Peninsular Malaysia, which has been described to be extinct for the past few decades. Through the means of both morphometric and molecular approaches, the species of *Tenuulosa* species in Perak has been identified as *T. ilisha*. This is the first recorded study to morphologically and genetically authenticate the *Tenuulosa* species in Perak in Malaysia. Though, this study has found that there are variations in morphometric and meristic characters. The morphological variations suggest the possibility that the samples are divided into two groups with the second group is the later localized generation of the first group. In addition, the genetic relationship of Malaysia *T. ilisha* is closest to that of Thailand, followed with the Indian and West Africa. These raise other research questions. Therefore, further study should be done to understand more about the biology and life history of Malaysian *T. ilisha* that inter-related with its genetic characteristics. Such studies are important to prevent this invaluable species from disappearing again in the future.

Acknowledgements. The study was supported by Universiti Malaysia Terengganu under GGP grant (68007/2013/102) and Ministry of Higher Education under FRGS grant 59245. Appreciation goes Biosystem Laboratory, School of Fisheries and Aquaculture Sciences and Institute of Tropical Aquaculture (AKUATROP) and Mrs. Siti Tafzilmeriam Sheikh Abdul Kadir, South China Sea Repository & Reference Center, Institute of Oceanography and Environment (INOS) for the great assistance given in conducting the study.

References

- Abdul Aziz A. H., Zakirah T., Nabilah M. A., Nur Asma A., Danish Daniel M. A., Lilian W., Seah Y. G., Mat Jaafar T. N. A., Ambok Bolong A. M., Kasim A. A. A., Shahreza M. S., 2015 Mitochondrial DNA diversity of Terubok (*Tenuulosa toli*) from Daro and Mukah, Sarawak inferred by partial cytochrome b (Cyt-B). *Journal of Fisheries and Aquatic Science* 10:92-101.
- Altschul S. F., Gish W., Miller W., Myers E. W., Lipman D. J., 1990 Basic local alignment search tool. *Journal of Molecular Biology* 215(3): 403-410.
- Arai T., Amalina R., 2014 New record of a tropical shad *Tenuulosa ilisha* (Teleostei: Clupeidae) in Malaysian waters. *Marine Biodiversity Records* 7(e66): 1-4.
- Blaber S. J. M., Milton D. A., Pang J., Wong P., Boon-Teck O., Nyigo L., Lubim D., 1996 The life history of the tropical shad *Tenuulosa toli* from Sarawak: first evidence of the protandry in the Clupeiformes? *Environmental Biology of Fishes* 46(3):225-242.
- Darlina M. N., Masazurah A. R., Jayasankar P., Jamsari A. F. J., Siti A. M. N., 2011 Morphometric and molecular analysis of mackerel (*Rastrelliger* spp) from the west coast of Peninsular Malaysia. *Genetics and Molecular Research* 10(3):2078-2092.
- De Silva M., Liyanage N., 2009 Morphological variation of *Puntius bimaculatus* with respect to altitudinal differences and five major river basins of Sri Lanka. *Ruhuna Journal of Science* 4:51-64.
- Gambang A. C., 1988 Some aspects of the biology of the Ikan Terubok *Hilsa toli* (Valenciennes) (Family: Clupeidae) at the lower Batang Lupar river, Sarawak. Department of Fisheries, Department of Agriculture, Malaysia. *Fisheries Buletin* 53:1-22.
- Garcia G., Rios N., Gutierrez V., Varela J. G., Fernandez C. B., Pardo B. G., Portela P. M., 2014 Promiscuous speciation with gene flow in silverside fish genus *Odontesthes* (Atheriniformes, Atherinopsidae) from south western Atlantic Ocean basins. *PLoS ONE* 9(8):e104659.

- Haniffa M. A., Abiya S. J., Milton J., Ramesh K., Bhat A. A., Chelliah A., 2014 Morphometric, meristic and ISSR marker systems for species identification and evolutionary analysis in five Indian Channids. *Biochemical Systematics and Ecology* 55: 131-136.
- Hanzawa N., Gotoh R. O., Sekimoto H., Goto T. V., Chiba S. N., Kuriwa K., Tamate H. B., 2012 Genetic diversity and evolution of marine animals isolated in marine lakes. In: *Analysis of genetic variation in animals*. InTech, Croatia, pp. 121-150.
- Hong X., Zhu X., Chen K., Pan D., Li K., 2013 Morphometric analysis of *Tenualosa ilisha*, *Alosa sapidissima* and *Tenualosa reevesii*. *Journal of South China Agricultural University* 34(2):203-206.
- Hubbs C. L., Lagler K. F., 1964 *Fishes of the Great Lakes region*. The University of Michigan Press, Ann Arbor, 213 pp.
- Jannatul-Fariyah Z., Nur Asma A., Shahreza M. S., 2011 Molecular characteristics of partial cytochrome C oxidase subunit 1 (*CO1*) on *Tenualosa toli* from Sarawak. UMT 11th International Annual Symposium on Sustainability Science and Management 11:91-96.
- Jorgensen H. B. H., Pertoldi C., Hansen M. M., Ruzzante D. E., Loeschcke V., 2008 Genetic and environmental correlates of morphological variation in a marine fish: the case of Baltic Sea herring (*Clupea harengus*). *Canadian Journal of Fisheries and Aquatic Sciences* 65:389-400.
- Kartavtsev Y. P., Lee J. S., 2006 Analysis of nucleotide diversity at the Cytochrome b and Cytochrome Oxidase 1 genes at the population, species, and genus levels. *Russian Journal of Genetics* 42(4):341-362.
- Kelley J. L., Passow C. N., Plath M., Rodriguez L. A., Yee M. C., Tobler M., 2012 Genomic resources for a model in adaptation and speciation research: characterization of the *Poecilia mexicana* transcriptome. *BMC Genomics* 13(652):1-13.
- Kolangi-Miandare H., Askari G., Fadakar D., Aghilnegad M., Siti-Azizah M. N., 2013 The biometric and Cytochrome Oxidase subunit I (*CO1*) gene sequence analysis of *Syngnathus abaster* (Teleostei: Syngnathidae) in Caspian Sea. *Molecular Biology Research Communications* 2(4):133-142.
- Li Y. L., Kong X. Y., Yu Z. N., Kong J., Ma S., Chen L. M., 2009 Genetic diversity and historical demography of Chinese shrimp *Feneropenaeus chinensis* in Yellow Sea and Bohai Sea based on mitochondrial DNA analysis. *African Journal of Biotechnology* 8(7):1193-1202.
- McDowall R. M., 1988 *Diadromy in fishes: migrations between freshwater and marine environments*. Croom Helm, London, 308 pp.
- Motamedi M., Madjdzadeh S. M., Teimori A., Esmaeili H. R., Mohsenzadeh S., 2014 Morphological and molecular perspective on geographical differentiation of *Barbus* populations (Actinopterygii; Cyprinidae) within Iranian freshwater drainages. *Turkish Journal of Fisheries and Aquatic Sciences* 14:339-351.
- Munroe T. A., Wongratana T., Nizinski M. S., 1999 Clupeidae. Herrings (also, sardines, shads, sprats, pilchards, and menhadens). In: *FAO species identification guide for fishery purposes. The living marine resources of Western Central Pacific. Volume 3. Batoid fishes, chimeras and bony fishes part1 (Elopidae to Liophryidae)*. Carpenter K. E., Niem V. H. (eds), FAO Rome, pp. 1775-1821.
- Narejo N. T., Lashari P. K., Jafri S. I. H., 2008 Morphometric and meristic differences between two types of Palla, *Tenualosa ilisha* (Hamilton) from River Indus, Pakistan. *Pakistan Journal of Zoology* 40(1):31-35.
- Palumbi S. R., 1996 *Nucleic acids II: the polymerase chain reaction*. In: *Molecular systematics*. Hillis D. M., Moritz C., Mable B. K. (eds), Sinauer Sunderland: Mass, pp. 205-221.
- Phillip W. P. H., 2001 The status of Terubok (*Tenualosa toli*) fishery in Sarawak. *Proceedings of the International Terubok Conference Sarawak, Malaysia*, Sarawak Development Institutes 1:91-99.
- Pillay S. R., Rosa H., 1963 Synopsis of biological data on hilsa, *Hilsa hilsa* (Hamilton). *FAO Fisheries Biology Synopsis* 25:1-60.
- Plath M., Hermann B., Schroder C., Riesch R., Tobler M., Garcia de Leon F. J., Schlupp I., Tiedemann R., 2010 Locally adapted fish populations maintain small-scale genetic differentiation despite perturbation by catastrophic flood event. *BMC Evolutionary Biology* 10:256.

- Rahim K. A. A., Tumiran M. H., Long S. M., Esa Y., Husaini A. A. S. A., 2014 The status of *Tenuulosa toli* (Valenciennes, 1847) in the south-west coast of Sarawak, Malaysia. *Kuroshio Science* 8(1):39-44.
- Seah Y. G., Abdullah S., Zaidi C. C., Mazlan A. G., 2009 Systematic accounts and some aspects of feeding and reproductive biology of ponyfishes (Perciformes: Leiognathidae). *Sains Malaysiana* 38(1):47-56.
- Seah Y. G., Mazlan A. G., Abdullah S., Zaidi C. C., Usup G., Mohamed C. A. R., 2011 Feeding guild of the dominant trawl species in the southeastern waters of Peninsular Malaysia. *Journal of Biological Sciences* 11(2):221-225.
- Siti-Balkhis A. B., Jamsari A. F. J., Hwai T. S., Yasin Z., Siti-Azizah M. N., 2011 Evidence of geographical structuring in the Malaysian snakehead, *Channa striata* based on partial segment of the *CO1* gene. *Genetics and Molecular Biology* 34(3):520-523.
- Stelbrink B., Stoger I., Hadiaty R. K., Schliewen U. K., Herder F., 2014 Age estimates for an adaptive lake fish radiation, its mitochondrial introgression, and an unexpected sister group: sailfin silversides of the Malili Lakes system in Sulawesi. *BMC Evolutionary Biology* 14(94):1-14.
- Song L. M., Munian K., Rashid Z. A., Bhasu S., 2013 Characterisation of Asian snakehead murrel *Channa striata* (Channidae) in Malaysia: an insight into molecular data and morphological approach. *The Scientific World Journal* 2013: 1-16.
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M., Kumar S., 2011 MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28(10):2731-2739.
- Tinti F., Ungaro N., Pasolini P., De Panfilis M., Garoia F., Guarniero I., Sabelli B., Marano G., Piccinetti C., 2003 Development of molecular and morphological markers to improve species-specific monitoring and systematics of north-east Atlantic and Mediterranean skates (Rajiformes). *Journal of Experimental Marine Biology and Ecology* 288:149-165.
- Turan C., Oral M., Ozturk B., Duzgunes E., 2006 Morphometric and meristic variations between stocks of bluefish (*Pomatomus saltatrix*) in the Black, Marmara, Aegean and northeastern Mediterranean Seas. *Fisheries Research* 79:139-147.
- Watanabe S., Aoyama J., Tsukamoto K., 2008 The use of morphological and molecular genetic variations to evaluate subspecies issues in the genus *Anguilla*. *Coastal Marine Science* 32(1):19-29.
- Whitehead P. J. P., 1985 FAO species catalogue. Vol. 7. Clupeoid fishes of the world (suborder Clupeoidei): An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, shads, anchovies, and wolffherrings. Part 1 - Chirocentridae, Clupeidae and Pristigasteridae. *FAO Fisheries Synopsis* 125(7):303.

Received: 01 November 2016. Accepted: 10 December 2016. Published online: 27 December 2016.

Authors:

Nurul Nadira Arjunaidi, School of Fisheries and Aquaculture Sciences, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Malaysia, e-mail: nadiraarjunaidi@gmail.com

Muhammad Faiz Zakaria, School of Marine Science and Environment, 21030 Kuala Terengganu, Malaysia, e-mail: faizzakaria92@gmail.com

Abdul Hadi Abdul Aziz, School of Fisheries and Aquaculture Sciences, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Malaysia, e-mail: hadiaziz41@gmail.com

Md. Sheriff Shahreza, School of Fisheries and Aquaculture Sciences & Institute of Tropical Aquaculture, 21030 Kuala Terengganu, Malaysia, email: shahreza@umt.edu.my

Tun Nurul Aimi Mat Jaafar, School of Fisheries and Aquaculture Sciences & Institute of Tropical Aquaculture, 21030 Kuala Terengganu, Malaysia, e-mail: tun_aimi@umt.edu.my

Seah Ying Giat, School of Fisheries and Aquaculture Sciences & Fish Division South China Sea Repository and Reference Center, Institute of Oceanography and Environment, 21030 Kuala Terengganu, Malaysia, e-mail: ygseah@umt.edu.my

Nur Asma Ariffin, School of Fisheries and Aquaculture Sciences & Institute of Tropical Aquaculture, 21030 Kuala Terengganu, Malaysia, e-mail: nurasma@umt.edu.my

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Arjunaidi N. N., Zakaria M. F., Abdul Aziz A. H., Shahreza M. S., Mat Jaafar T. N. A., Seah Y. G., Nur Asma A., 2016 Authentication of *Tenuulosa* species in Perak River, Malaysia: application of morphological measurement and molecular analysis of partial *CO1* and 16S genes to resolve species ambiguity. *AAFL Bioflux* 9(6):1355-1363.