



Complexity of carbon sources and the impact on biofloc integrity and quality in tilapia (*Oreochromis niloticus*) tanks

¹Osama M. El-Husseiny, ²Ashraf M. A.-S. Goda, ¹Rania S. Mabroke, ²Mohamed Soaudy

¹ Faculty of Agriculture, Cairo University, Giza, Egypt; ² Fish Nutrition Laboratory, National Institute of Oceanography and Fisheries (NIOF), Cairo, Egypt. Corresponding author: R. S. Mabroke, Rania.Ahmed@agr.cu.edu.eg; ranciasms2002@yahoo.com

Abstract. Organic carbon sources are the key factor in optimizing the biofloc system conditions. Our concern was to study the effect of different carbon sources on biofloc conditions and tilapia performance. Biofloc treatments with five different organic carbon simple sources (glucose, molasses) and complex sources (starch, wheat bran, cellulose) were conducted in presence of control (clear water). No significant differences ($p > 0.05$) were noticed among different organic carbon sources regarding tilapia performance. Complex carbon source represented in wheat bran and cellulose showed less fluctuation in the values of NH_4^+ and NO_2 during the experimental period than the other carbon sources. The precipitated biofloc from both wheat bran and cellulose showed the highest fat content (8.08 and 7.72 respectively). In terms of heterotrophic bacteria production, plankton count and biofloc nutritional content, cellulose seems to be the better choice. From the nutritional and economical points of view, using agriculture by product with high cellulose content as a carbon source in biofloc system is more reasonable, guarantee positive environmental impact and aquaculture sustainability.

Key Words: organic carbon sources, biofloc, Nile tilapia, growth performance.

Introduction. With more expansion of biofloc system application around the world, optimizing the conditions for biofloc system is needed for efficient water utilization and for optimal fish growth performance. These efforts are especially needed under conditions of water shortages and rise in world population. Optimal biofloc conditions guarantee more aquaculture intensification, best profiteering of land, water and feed recourses. Intensification of fish culture causes waste accumulation in forms of solids (e.g. feces and uneaten feed) and nutrients (e.g. nitrogen and phosphorus) which have adverse effect on the environment. Biofloc is considered a waste treatment system under condition of zero water exchange. Carbon sources supplementation are the backbone for initiating the system, where affording Carbon: Nitrogen ratio (C:N ratio) of 10 guarantee the activation of the heterotrophic bacteria (Avnimelech 1999; De Schryver et al 2008). The last have the ability to assimilate the nitrogen wastes and recycle it into microbial protein which considered as secondary feed source for species such as tilapia, carp and shrimp. Consequently, water quality and performance of fish culture could be enhanced (Avnimelech 2009; Crab et al 2010). Previous research work examined carbon sources for accessibility, dissolving rate of carbohydrate in water, bioavailability and costs under biofloc condition, especially for shrimp culture. Molasses, corn flour and wheat bran mixture (3:1:1, respectively) improved shrimp (*Litopenaeus vannamei*) growth performance and floc nutritional value (Wang et al 2016). Rice flower showed better values than molasses regarding growth and immune response of black tiger shrimp *Penaeus monodon* cultured under biofloc system Kumar et al (2017). Recently, Khanjani et al (2017) investigated the effect of molasses, starch, wheat flour and mixture of them on the performance of Pacific white shrimp *L. vannamei* where molasses given the best performance of shrimp. Crab et al (2010) developed biofloc system under different

carbon sources acetate, glycerol and glucose to determine the possible use of bioflocs as a feed for shrimp post larvae *Macrobrachium rosenbergii* where glycerol+bacillus was appropriate for shrimp culture. For our knowledge some studies evaluated the effect of supplementing biofloc system with carbon sources of different complexity on tilapia culture. The experiment was designed to determine the effect of organic carbon source with different complexity grade (glucose, molasses, starch, wheat bran, cellulose) on the maintaining of biofloc system condition, tilapia growth performance, water quality parameters, feed utilization, floc chemical composition and fish hematological parameters meanwhile detecting the most efficient and economic carbon sources for tilapia culture under biofloc system.

Material and Method

Fish stocking and rearing techniques. Fingerlings of Nile tilapia, *Oreochromis niloticus* were obtained from El-Fayum governorate and transferred to the farm of El-Kanater El-Khayria, National Institute of Oceanography and Fisheries (NIOF), Cairo. Fingerlings were fed on a diet with a crude protein of 30% and were acclimatized to laboratory condition for two weeks. Eighteen (18) cylindrical plastic tanks (with water capacity of 150 L and filled to 100 L) were stocked with fingerlings at a rate of 10 fish tank⁻¹ (100 fish m⁻³), with an initial weight of 14.00±0.03 g. The six experimental treatments were represented in triplicate. Tanks were supplemented with well water source and were continuously aerated using air pump (SIEMENS, 1LE1002-1BA22-2AA4, 2-pole * Size 112M, Czech Republic) using porous stone with length of 4.7 cm. As for the control (clear system), water was exchanged biweekly, while for experimental biofloc tanks, no water exchange was done (zero water exchange) except to compensate the evaporated water.

Diet preparation. A homogenous mixture was used to blend the dry ingredient of basal diet. The ration was then pelleted using the mill (California Pellet Mill, San Francisco, CA, USA), solar dried and stored at -4°C. Experimental fish were fed six days a week on the basal diet (26% protein) for 85 days (Table 1). The daily ration (3% of stocked biomass) was delivered three times a day (10.00, 12.00 and 14.00 hr) in equal portions. Fish were weighted every 15 days and daily feed ration was adjusted accordingly.

Table 1

Formulation and chemical composition of the basal diet

Ingredients	Diet formula
Fish meal	6
Soya bean meal	43
Yellow corn	20
Wheat bran	23
Soya oil	6
Vit. and Min. premix ¹	1.45
CMC ²	0.5
Vit. C	0.05
<i>Proximate composition</i>	
Dry matter (%)	89.70
Protein (%)	25.93
Lipid (%)	14.48
Total carbohydrate (%) ³	43.21
Ash (%)	6.08
Gross energy (MJ /kg) ⁴	19.26
Metabolizable energy (MJ /kg) ⁵	13.84
Protein energy (mg CP/ KJ DE)	74.28

¹ Vitamins and minerals mixture each 3 Kg of mixture contain: 12 m.IU vit A, 12 mIU vit D3, 10 g vit. E, 2 g vit. K, 1 g vit. B1, 5 g vit. B2, 1.5 g vit. B6, 10 mg vit. B12, 30 g niacin, 1000 mg folic acid, 50 mg biotin, 10 g banathonic acid, 50 g zinc, 30g iron, 60 g manganese, 10 g copper, 100 mg cobalt, 100 mg selenium, 1000 mg iodine; ² CMC: carboxy methyl cellulose; ³ Total carbohydrate = 100-(CP + EE + ash); ⁴ Calculated using gross calorific values of 23.63, 39.52 and 17.15 KJ g⁻¹ for protein, fat and carbohydrate, respectively according to Brett (1971); ⁵ Calculated metabolizable energy = 70% of determined GE value (Hepher et al 1983).

Managing of biofloc tanks. C:N ratio of 10:1 (De Schryver et al 2008) were maintained in different experimental tanks using different carbon sources: glucose, molasses, starch, wheat bran or cellulose that were examined for optimal biofloc condition. Daily addition of different carbon sources was applied. Carbon sources were completely mixed with water sample of the experimental tanks prior spreading into the whole water body.

Water quality assessment. Water temperature and dissolved oxygen (HI 9146-HANNA interment, USA) were recorded weekly. pH values were recorded twice a week (Orion pH meter, Abilene, Texas, USA). Ammonia, nitrite and nitrate were detected biweekly (APHA 1992). Total suspended solid (TSS) values were measured twice during the experimental period from June to September 2014 using (Multi direct Iovibond). Total bacterial count (TBC) was determined according to (AOAC 1995; APHA 1992) then reported as CFU mL⁻¹.

Chemical analysis. At experiment end, diet sample along with random samples of five individual fish was sampled from each tank of different treatment and precipitated flocs were collected from different treatments for determination of proximate composition. Fish and biofloc sample were dried in an oven at 80°C till constant weight then were grounded and stored at -20°C for subsequent analysis. By incineration at 550°C for 6 h, ash content was detected. Crude protein was determined by micro-Kjeldhal method, %N × 6.25 (using Kjeltach auto analyzer, Model 1030, Tecator, Höganäs, Sweden). Soxhlet extraction with diethyl ether (40-60°C) was used to estimate crude fat content of different samples, the chemical analysis determined according to AOAC (1995).

Hematological parameters. Tubs coated with 10% EDTA (ethylene diamine tetra acetate) were used to collect blood samples for estimation of hematocrit (Ht). The cyanomet hemoglobin method was used to detect the value of hemoglobin (Hb) and white blood cell (WBCs) was estimated by the method of Martins et al (2004).

Statistical analysis. One-way ANOVA was used to analyze experimental data and to detect the best carbon source under biofloc system using the SAS v9.0.0 (2004) program. Duncan test (1955) at $p \leq 0.05$ level of significant was applied.

Results and Discussion

Water quality parameters. Water temperature, pH, dissolved oxygen (DO), ammonia (NH₄⁺), nitrite (NO₂) and nitrate (NO₃) are shown in Table 2. No significant differences ($p > 0.05$) were observed for water temperature, pH, NH₄⁺, NO₂ and NO₃ values among different carbon sources during the experimental period. DO values had significant differences ($p < 0.05$) among different treatments. The fluctuations of NH₄⁺, NO₂ and NO₃ during experimental period are shown in Figure 1. Values of NH₄⁺ and NO₂ were high at the beginning of the experiment then decreased along with degradation of different carbon sources subsequently increase of bacterial count. The changes in NH₄⁺, NO₂ and NO₃ values were in the acceptable range for Nile tilapia (DeLong et al 2009). Generally, values of chemical water quality parameters decreased over time for all treatments. Supplementing experimental tanks with organic carbon sources enhanced the heterotrophic bacterial growth; subsequently increase of nitrogen fixation and decrease ammonia and nitrite which enhanced the water quality (Luo et al 2013, 2014). Despite no water discharged in biofloc treatments were done, no significant differences among control and biofloc treatments in different water quality parameters were found, these in consonance with Nootong & Pavasant (2011). Application of biofloc technique enhanced the water quality via minimizing the level of NH₄⁺, NO₂ in carp ponds by elevating the C:N ratio (Wang et al 2015). Same was recognized by Prajith & Madhusoodana (2011) for shrimp culture. Khanjani et al (2017) stated that there were no significant differences among molasses, starch, and wheat flour regarding water quality parameters ammonia, nitrite and nitrate in shrimp *L. vannamei* tanks under biofloc system. Molasses (simple) or rice flour (complex) carbon sources showed the same effects on water quality parameters in shrimp tanks that fed either 30 or 40% crude protein under condition of no water exchange (Kumar et al 2017). Biofloc treatments recorded significantly lower DO

than control, this may be due to (i) the decomposition of organic matter by microbial metabolism process which consumed oxygen from water (Khanjani et al 2017), (ii) increased the microbial load in biofloc tanks in presence of high C:N ratio.

Table 2
Water quality parameters of different treatments during the experimental period

Parameters	Control	Different carbon sources					±MSE	P value
		Glucose	Molasses	Starch	Bran	Cellulose		
Temperature(°C)	28.75	28.72	28.70	28.74	28.75	28.77	0.33	1.0000
DO (mg L ⁻¹)	5.71 ^a	4.90 ^b	4.61 ^b	4.81 ^b	4.86 ^b	4.96 ^b	0.25	0.0543
PH	8.58	8.25	8.31	8.39	8.30	8.32	0.14	0.6040
NH ₄ ⁺ (mg L ⁻¹)	0.96	1.21	1.12	1.22	1.15	0.95	0.34	0.9877
NO ₂ (mg L ⁻¹)	0.19	0.24	0.23	0.33	0.34	0.18	0.08	0.5517
NO ₃ (mg L ⁻¹)	0.76	1.18	1.22	1.10	1.53	1.15	0.17	0.0583
TSS	-	537 ^a	549 ^a	359 ^b	553 ^a	479 ^a	38.35	0.0074

Data are presented as means±standard error (SE). Means followed by different letters in each rows are significantly (p < 0.05) different.




Figure 1. Fluctuations of NH₄⁺, NO₂ and NO₃ during experimental period.

In general, total suspended solids (TSS) values increased gradually during the experimental period, and ranged from 359 to 553 mg L⁻¹ among treatment of different carbon sources. The highest TSS value was recorded for wheat bran while, the least value was noticed for starch treatment. Since the recommended level of TSS should not reach 500 mg L⁻¹ (Samocho et al 2007), precipitation of biofloc was done after 30 days of the experiment beginning and at the end. Even though, elevations of TSS over optimal level were recorded, thus a procedure of floc precipitation had to be done more frequently to avoid elevation of TSS levels.

Total bacterial counts (TBC) are presented in Figure 2. The TBC counts significantly differed ($p > 0.05$) among different carbon sources. Cellulose, bran and molasses treatments were recorded the highest total count, respectively, compared with the others. TBC count in simple carbon sources treatments were lower than in complex sources treatments. It could be suggested that the last degraded gradually and slowly which led to higher carbon concentration in late stages of the experimental time, subsequently bacterial count increased by the end of the experiment. In contrary, Kumar et al (2017) observed that molasses which is considered as a simple carbon source showed higher microbial load than a complex carbon source as rice flour, under condition of feeding fish on 40% dietary protein. He suggested that this could be due to molasses contain sucrose (disaccharides) in contrast with rice flour, where the last contains polysaccharide. Serra et al (2015) who reported that the speed of carbon degradation rate of simple carbohydrate (molasses and dextrose) is faster than the complex carbohydrate (rice bran), that provide carbon rapidly in water environment as a substrate of bacteria. The positive effect of different complex carbon sources on biofloc shrimp culture was reported by Prajith & Madhusoodana (2011) where the total number of heterotrophic bacteria increased in water (99.9, 74.45, 60.39, 88.7, and 72.9 x 10³, respectively), without significant differences among different carbon sources (tapioca flour, yam flour, wheat flour, rice flour and potato flour). Generally, slow degradation of complex carbon sources may have higher constant concentration of organic carbon during the experimental period, which gradually increased the count of heterotrophic bacteria. Subsequently, there was less fluctuation in values of NH₄⁺ and NO₃. The constantly of last parameters may affect positively the health condition of aquatic organisms.




Figure 2. The effect of different organic carbon sources on total bacterial count under biofloc system compared with control.

Growth performance and feed utilization of tilapia. The growth performance (final body weight (FBW), weight gain (WG) and specific growth rate (SGR)) and feed utilization (feed intake (FI), feed conversion ratio (FCR) and protein efficiency ratio (PER)) are given in Table 3. No significant differences were noticed ($p > 0.05$) among different carbon sources regarding fish performance. Generally, all organic carbon sources showed superiority in growth performance and feed utilization compared to the control. Addition

of different carbon sources revealed a balanced C:N ratio, subsequently activation of the heterotrophic bacteria. This along with micro organisms such as zooplankton protozoa and others acted as secondary feed source (Schneider et al 2005). Luo et al (2014) concluded that WG, SGR and FCR were better in biofloc system than recirculating aquaculture systems (RAS). Long et al (2015) noted the same results for tilapia. Furthermore, Ekasari et al (2015) demonstrated better productive performance of Nile tilapia larva under biofloc condition was gained. Crucian carp *Carassius auratus* showed improved growth performance with increasing of the C:N up to 25:1 (Wang et al 2015).

Table 3
Growth performances and feed utilization of tilapia under biofloc system with different carbon sources compared with control (clear system)

Parameters	Control	Different carbon sources					±MSE	P value
		Glucose	Molasses	Starch	Bran	Cellulose		
Initial weight	14.06	14.01	13.98	14.01	14.04	13.99	0.03	0.3071
Final weight	22.81	26.69	26.14	24.35	26.24	24.50	1.10	0.1277
Weight gain	7.20 ^b	10.20 ^a	11.63 ^a	10.30 ^a	12.20 ^a	10.00 ^a	0.83	0.0211
SGR (% day)	0.57	0.75	0.73	0.65	0.73	0.66	0.05	0.1234
Feed intake	35.35	36.65	37.85	38.59	40.18	37.48	0.86	0.5034
FCR	5.60 ^a	4.27 ^b	3.50 ^b	3.80 ^b	3.40 ^b	3.90 ^b	0.32	0.0118
PER	0.70 ^b	0.91 ^{ab}	1.14 ^a	1.02 ^a	1.17 ^a	0.99 ^a	0.08	0.0248

Data are presented as means±standard error (SE). Means followed by different letters in each rows are significantly ($p < 0.05$) different. WG = Final body weight (g) - Initial body weight (g); SGR% = $(\ln \text{FBW} - \ln \text{IBW}) / t \times 100$; FCR = Diet intake (g)/weight gain (g); PER = weight gain (g)/protein intake (g).

Growth performance parameters didn't clearly reflect the effect of different carbon sources under biofloc system condition. Kumar et al (2017) suggested that the complex carbon source rice flour positively affected shrimp (*P. monodon*) growth performance than simple carbon source molasses. Same trend was recorded by Serra et al (2015) who reported that the best result of shrimp (*L. vannamei*) growth recorded for rice bran treatment as a complex carbohydrate source in comparison with dextrose as a simple carbon source. Many authors suggested no differences among organic carbon sources as a control element in biofloc system were noticed (Prajith & Madhusoodana 2011; Khanjani et al 2017). Then, choosing cheaper carbon sources especially complex sources such as agriculture by product showed to be more economical and eco-friendly.

The poor growth performance and feed utilization that recorded under either control or floc treatments could be due to; low diet stability because of using roughness ingredient, spawning that showed under both floc and control treatments. The same most of reasons were recorded by Azim & Little (2008). Furthermore, elevation of TSS in biofloc tanks may be another reason behind the poor growth performance. It seems that under condition of biofloc, the settling process should be preceded more frequently during the experiment duration. Adverse effect on shrimp performance was recognized by Furtado et al (2011) with the excess of TSS values. Elevation of TSS concentration above recommended level (500 mg L^{-1}) resulted in adverse effect on growth performance and production of tilapia fish (Azim & Little 2008). Low growth performance and feed utilization in present case could be attributed to high values of TSS which exceeded the optimal range in biofloc treatments.

Chemical composition of fish and biofloc. Carcass chemical compositions of tilapia are presented in Table 4. No significant differences ($p > 0.05$) among treatments regarding ash, fat and protein contents were noticed. Significant differences ($p < 0.05$) were showed in dry matter, where the highest values were recorded for wheat bran. Khanjani et al (2017) noticed higher fat content of shrimp *L. vannamei* in complex carbon source treatment (wheat flour) compared to treatments of molasses, starch or their mixture. The highest fat content of tilapia in our results showed in wheat bran as a complex carbon source without significant difference with other treatments. Long et al (2015) found that lipid content increased in fish carcass under the biofloc treatment

compared with the control, while no significant differences in crude protein and ash content were recorded. Luo et al (2014) showed no significant differences in protein and fat composition of tilapia muscle between either RAS or biofloc system.

Table 4
Chemical composition of fish under biofloc system with different carbon sources compared with control (clear system)

Parameters	Control	Different carbon sources					±MSE	P value
		Glucose	Molasses	Starch	Bran	Cellulose		
Dry matter	23.11 ^{bc}	23.24 ^{bc}	24.07 ^{ab}	22.50 ^c	25.08 ^a	24.16 ^{ab}	0.43	0.0428
Ash	19.13	19.12	19.38	20.62	19.24	20.10	0.62	0.5818
Fat	19.83	18.68	19.46	17.34	20.14	19.80	0.66	0.1155
Protein	59.48	59.28	57.7	60.17	57.86	59.21	1.04	0.6892

Data are presented as means±standard error (SE). Means followed by different letters in each rows are significantly (p < 0.05) different.

Biofloc chemical composition under different carbon sources are tabulated in Table 5. The heights protein and fat content listed for starch and wheat bran treatment, respectively. Crude protein, lipid, ash and carbohydrate values of previous studies ranged between 18.4-49%; 0.1-4%; 11.8-42% and 19-36.4%, respectively (Azim & Little 2008; Kuhn et al 2010, 2016; Emerenciano et al 2013; Maica et al 2012; Dantas et al 2016). The resulted values in our study lay in the same ranges except for fat content where showed higher values that reached 8.08%. Carbon sources may reflect the biochemical composition of biofloc differently. Same was suggested by Crab et al (2010) where different carbon sources (glycerol+bacillus, glucose and acetate) resulted in different chemical composition of biofloc. Wheat flour resulted in the highest protein and lipid contents (30.73, 2.18, respectively) in the study of Khanjani et al (2017).

Table 5
Chemical composition of biofloc with different carbon source

Parameters	Different carbon sources					±MSE	P value
	Glucose	Molasses	Starch	Bran	Cellulose		
Protein	28.34 ^{bc}	32.74 ^{ab}	33.70 ^a	27.07 ^c	29.10 ^{abc}	1.56	0.0482
Ash	19.27	20.97	21.59	19.21	20.75	0.91	0.1885
Fat	3.93 ^c	5.21 ^{bc}	6.09 ^{abc}	8.08 ^a	7.72 ^{ab}	0.86	0.0239

Data are presented as means±standard error (SE). Means followed by different letters in each rows are significantly (p < 0.05) different.

Hematological blood parameters of fish. Hematological parameters including: hemoglobin (HB), red blood cell (RBC), hematocrit (HTC), and white blood cell (WBC) of tilapia under different organic carbon sources compared with the control are presented in Table 6.

Table 6
Hematological parameters of fish with different carbon sources in biofloc system compare with control (clear system)

Parameters	Control	Different carbon sources					±MSE	P value
		Glucose	Molasses	Starch	Bran	Cellulose		
HGB (g dl)	4.67	4.4	5.03	4.17	5.47	5.13	0.62	0.6873
HCT (%)	19.03	18.93	22.67	17.6	23.97	20.77	2.97	0.3550
RBC (10 ⁶ /ul)	1.36	1.56	1.71	1.24	1.76	1.7	0.19	0.3550
WBC (10 ³ /ul)	203.04	266.46	269.11	263.81	365.63	351.34	58.25	0.4184

Data are presented as means±standard error (SE). Means followed by different letters in each rows are significantly (p < 0.05) different.

Despite no significant differences ($p > 0.05$) among treatments in all blood parameters were recorded, the highest WBC values were recorded for complex carbon source (wheat bran and cellulose) treatments. The same trend was observed for other blood parameters. It seems that biofloc condition didn't affect fish health negatively compared with the clear water condition represented by the control. The same was reported by Long et al (2015) who suggested that no significant effect of biofloc up on blood hematology parameters, especially on RBC, WBC, Hb, and HTC. Furthermore, Azim & Little (2008) suggested the same range of hematocrit value for tilapia under either biofloc treatment (BFT) or recirculating aquaculture system (RAS).

Conclusions. Biofloc system with different carbon sources achieved superiority over control system in all parameters. There were no significant differences among biofloc treatments regarding tilapia performance. The slow degradation of complex carbon source (wheat bran and cellulose) delayed the improvement of water quality at the beginning of the experiment with less fluctuation values of NH_4^+ and NO_3^- . The present study suggested that complex carbon sources especially cellulose were proper choice interims of bacterial count, microorganisms diversity and quality of biofloc composition. From the nutritional and economical points of view choosing cheap agriculture by product with high cellulose content as a source of carbon in biofloc system is reasonable, may be resulted in positive impact on the environment and sustainability.

References

- American Public Health Association (APHA), 1992 Standard methods for the examination of water and waste water. 18th edition, American Public Health Association, Washington DC., 1268 pp.
- AOAC, 1995 Official methods of analysis. 15th edition, Association of Official Analytical Chemists, Arlington, VA, pp. 1-45.
- Avnimelech Y., 1999 Carbon/nitrogen ratio as a control element in aquaculture system. *Aquaculture* 176:227-235.
- Avnimelech Y., 2009 Biofloc technology - a practical guide book. The World Aquaculture Society, Baton Rouge, Louisiana, USA, 182 pp.
- Azim M. E., Little D. C., 2008 The biofloc technology (BFT) in indoor tanks: water quality, biofloc composition, and growth and welfare of Nile tilapia (*Oreochromis niloticus*). *Aquaculture* 283:29-35.
- Brett J. R., 1971 Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and fresh water ecology of sockeye salmon (*Oncorhynchus nerka*). *American Zoologist* 11:99-113.
- Crab R., Chielens B., Wille M., Bossier P., Verstraete W., 2010 The effect of different carbon sources on the nutritional value of bioflocs, a feed for *Macrobrachium rosenbergii* postlarvae. *Aquaculture Research* 41(4):559-567.
- Dantas E. M., Valle B. C. S., Brito C. M. S., Calazans N. K. F., Peixoto S. R. M., Soares R. B., 2016 Partial replacement of fishmeal with biofloc meal in the diet of post larvae of the Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture Nutrition* 22:335-342.
- De Schryver P., Crab R., Defoirdt T., Boon N., Verstraete W., 2008 The basics of bioflocs technology: the added value for aquaculture. *Aquaculture* 277:125-137.
- Delong D. P., Losordo T. M., Rakocy J. E., 2009 Tank culture of tilapia. SRAC publication 282, Southern Regional Aquaculture Center, 8 pp.
- Duncan D. B., 1955 Multiple range and multiple F tests. *Biometrics* 11(1):1-42.
- Ekasari J., Rivandi D. R., Firdausi A. P., Surawidjaja E. H., Zairin Jr. M., Bossier P., De Schryver P., 2015 Biofloc technology positively affects Nile tilapia (*Oreochromis niloticus*) larvae performance. *Aquaculture* 441:72-77.
- Emerenciano M., Cuzon G., Arévalo M., Miquelajaugui M. M., Gaxiola G., 2013 Effect of short-term fresh food supplementation on reproductive performance, biochemical composition, and fatty acid profile of *Litopenaeus vannamei* (Boone) reared under biofloc conditions. *Aquaculture International* 21(5):987-1007.

- Furtado P. S., Poersch L. H., Wasielesky W., 2011 Effect of calcium hydroxide, carbonate and sodium bicarbonate on water quality and zootechnical performance of shrimp *Litopenaeus vannamei* reared in bio-flocs technology (BFT) systems. *Aquaculture* 321(1-2):130-135.
- Hepher B., Liao I. C., Cheng S. H., Hsieh C. S., 1983 Food utilization by red tilapia. Effects of diet composition, feeding level and temperature on utilization efficiency for maintenance and growth. *Aquaculture* 32:255-275.
- Khanjani M. H., Sajjadi M. M., Alizadeh M., Sourinejad I., 2017 Nursery performance of Pacific white shrimp (*Litopenaeus vannamei* Boone, 1931) cultivated in a biofloc system: the effect of adding different carbon sources. *Aquaculture Research* 48(4):1491-1501.
- Kuhn D. D., Lawrence A. L., Boardman G. D., Patnaik S., Marsh L., Flick Jr. G. J., 2010 Evaluation of two types of bioflocs derived from biological treatment of fish effluent as feed ingredients for Pacific white shrimp, *Litopenaeus vannamei*. *Aquaculture* 303(1-4):28-33.
- Kuhn D. D., Lawrence A. L., Crockett J., Taylor D. P., 2016 Evaluation of bioflocs derived from confectionary food effluent water as a replacement feed ingredient for fishmeal or soy meal for shrimp. *Aquaculture* 454:66-71.
- Kumar S., Anand P. S. S., De D., Deo A. D., Ghoshal T. K., Sundaray J. K., Ponniah A. G., Jithendran K. P., Raja R. A., Biswas G., Lalitha N., 2017 Effects of biofloc under different carbon sources and protein levels on water quality, growth performance and immune responses in black tiger shrimp *Penaeus monodon* (Fabricius, 1978). *Aquaculture Research* 48(3):1168-1182.
- Long L., Yang J., Lib Y., Guan C., Wu F., 2015 Effect of biofloc technology on growth, digestive enzyme activity, hematology, and immune response of genetically improved farmed tilapia (*Oreochromis niloticus*). *Aquaculture* 448:135-141.
- Luo G. Z., Avnimelech Y., Pan Y. F., Tan H. X., 2013 Inorganic nitrogen dynamics in sequencing batch reactors using bioflocs technology to treat aquaculture sludge. *Aquacultural Engineering* 52:73-79.
- Luo G. Z., Gao Q., Wang C., Liu W., Sun D., Li L., Tan H., 2014 Growth, digestive activity, welfare, and partial cost-effectiveness of genetically improved farmed tilapia (*Oreochromis niloticus*) cultured in a recirculating aquaculture system and an indoor biofloc system. *Aquaculture* 422-423:1-7.
- Maicá P. F., de Borba M. R., Wasielesky Jr. W., 2012 Effect of low salinity on microbial floc composition and performance of *Litopenaeus vannamei* (Boone) juveniles reared in a zero-water-exchange super-intensive system. *Aquaculture Research* 43(3):361-370.
- Martins M. L., Nomura D. T., Myiazaki D. M. Y., Pilarsky F., Ribeiro K., de Castro M. P., de Campos C. F. M., 2004 Physiological and hematological response of *Oreochromis niloticus* (Osteichthyes: Cichlidae) exposed to single and consecutive stress of capture. *Acta Scientiarum. Animal Sciences* 26:449-456.
- Nootong K., Pavasant P., 2011 Effects of organic carbon addition in controlling inorganic nitrogen concentrations in a biofloc system. *Journal of the World Aquaculture Society* 42(3):339-346.
- Prajith K.K., Madhusoodana K.B., 2011 Application of Biofloc Technology (BFT) in the nursery rearing and farming of giant freshwater prawn, *Macrobrachium rosenbergii* (deMan) (Doctoral dissertation, Cochin University of Science and Technology).
- Samocha T. M., Patnaik S., Speed M., Ali A. M., Burger J. M., Almeida R. V., Ayub Z., Harisanto M., Horowitz A., Brook D. L., 2007 Use of molasses as carbon source in limited discharge nursery and grow-out systems for *Litopenaeus vannamei*. *Aquacultural Engineering* 36:184-191.
- SAS Institute Inc., 2004 INSIGHT® 9.1 User's Guide, Volumes 1 and 2. Cary, NC: SAS Institute Inc., 816 pp.
- Schneider O., Sereti V., Eding E. H., Verreth J. A. J., 2005 Analysis of nutrient flows in integrated intensive aquaculture systems. *Aquacultural Engineering* 32:379-401.

- Serra F. P., Gaona C. A. P., Furtado P. S., Poersch L. H., Wasielesky W., 2015 Use of different carbon sources for the biofloc system adopted during the nursery and grow-out culture of *Litopenaeus vannamei*. *Aquaculture International* 23(6):1325-1339.
- Wang C., Pan L., Zhang K., Xu W., Zhao D., Mei L., 2016 Effects of different carbon sources addition on nutrition composition and extracellular enzymes activity of bioflocs, and digestive enzymes activity and growth performance of *Litopenaeus vannamei* in zero-exchange culture tanks. *Aquaculture Research* 47(10):3307-3318.
- Wang G., Yu E., Xie J., Yu D., Li Z., Luo W., Qiu L., Zheng Z., 2015 Effect of C/N ratio on water quality in zero-water exchange tanks and the biofloc supplementation in feed on the growth performance of crucian carp, *Carassius auratus*. *Aquaculture* 443:98-104.

Received: 15 April 2018. Accepted: 10 June 2018. Published online: 21 June 2018.

Authors:

Osama Mohamed El-Husseiny, Faculty of Agriculture, Cairo University, El-Gamma St., 12613 Giza, Egypt, e-mail: osama.elhussiny@yahoo.com

Ashraf Mohamed Abd-Elsamee Goda, Fish Nutrition Laboratory, National Institute of Oceanography and Fisheries (NIOF), 101 Kasr ELAiny St., 11516 Cairo, Egypt, e-mail: goda_ashraf@yahoo.com

Rania Said Mabroke, Faculty of Agriculture, Cairo University, El-Gamma St., 12613 Giza, Egypt, e-mail: rania.ahmed@agr.cu.edu.eg

Mohamed Soaudy, Fish Nutrition Laboratory, National Institute of Oceanography and Fisheries (NIOF), 101 Kasr ELAiny St., 11516 Cairo, Egypt, e-mail: mrssouady@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

El-Husseiny O. M., Goda A. M. A. S., Mabroke R. S., Soaudy M., 2018 Complexity of carbon sources and the impact on biofloc integrity and quality in tilapia (*Oreochromis niloticus*) tanks. *AAFL Bioflux* 11(3):846-855.