



The economic value of the resource utilization of wetlands: comparative study of *beje* fisheries in North Hulu, Sungai Regency, South Kalimantan, Indonesia

¹Leila A. Sofia, ²Susi Nurlianti

¹ Study Program of Fisheries Agribusiness, Faculty of Fisheries and Marine, Lambung Mangkurat University, South Kalimantan, Indonesia; ² Graduate of Program of Fisheries Agribusiness, Faculty of Fisheries and Marine, Lambung Mangkurat University, South Kalimantan, Indonesia. Corresponding author: L. A. Sofia, leila.ariyani@ulm.ac.id

Abstract. *Beje* fishery in wetlands is generally managed by fishermen only on the basis of their own capital and experience, so that land is cleared as far as possible without considering the financial feasibility aspects of the business and the conservation of wetlands. The aims of the study were to determine the volume and value of production, and the feasibility of *beje* fishery business based on the number of management units. The location of the study was determined purposively, namely Papuyu River Village, North Hulu Sungai Regency of South Kalimantan, and samples were taken by census of 20 fisherman *beje* households. The samples are grouped into 3 groups based on the number of managed *beje* units are small groups (1-5 units); middle group (6-10 units); and large groups (more than 10 units). Primary data were analyzed using descriptive analysis and business feasibility analysis through investment criteria approach. The results showed that the highest total *beje* production volume per unit is categorized as the moderate level with production reaching 100.91 kg year⁻¹ with the highest production value per unit was approximately Rp 1,244,090 year⁻¹. The middle management group has better land use, capital and operational efficiency than small and large groups.

Key Words: inland fisheries, wetlands, conservation, feasibility studies, management.

Introduction. Wetlands cover nearly 6% of the earth's surface, including swamp, fen, peat lands or peat waters, either natural or artificial, permanent or transient, in the form of static or flowing freshwater, brackish or saline, including the area of sea water with receding depth not exceeding six meters (Stuip et al 2002). Wetland systems have benefited directly, or indirectly, to the world's population. A single wetland system may provide multiple types of ecosystem services depending on wetland conditions of type, location, condition, utilization, etc. (Whiteoak & Binney 2012). Wetlands play an important role in slowing down and storing flood waters (Leschine et al 1997), controlling pollution, contributing to local and national economies by producing resources and providing recreation (EPA 2006; Früh et al 2013; Das et al 2015).

Fisheries and aquaculture in the mainland has contributed more than 40% of the world production of finned fish were reported from at least 0.01% of the total volume of water on earth. This fishery provides food for billions and livelihoods for millions of the world's population (Lynch et al 2016). Traditional fishing is an important livelihood for most households (especially the poor) in wetland areas, both as a source of household protein as well as investment resources, business and cash income (David et al 2008). Similarly aquaculture in wetlands is able to provide fish production (nutrient supply) and high economic benefits for farmers (Chandra et al 2010; Olaoye et al 2014).

However, government policies in some countries still do not prioritize the use of wetlands for fisheries development, where the contribution of the economic value of the utilization of wetlands from fisheries is still considered lower than the contribution from

the use of wetlands for other economic activities (Cooke et al 2016). Many wetland areas are used massively for economic activities outside of fisheries, such as drainage for agriculture, settlement and development of areas, waste disposal sites, etc. that threaten the ecology of wetlands and the existence of fisheries resources. Ramsar data shows that in 1994 about 84% of registered wetlands had experienced or were threatened by ecological changes (Stuip et al 2002).

One of the many traditional fisheries systems developed in several areas of Indonesia's wetlands (such as Sumatra and Kalimantan) is *beje*. *Beje* fishery has been developed by the people of South Kalimantan from generation to generation, especially in the freshwater swamp area of North Hulu Sungai Regency. *Beje* is a pond in a deliberately created swampland and serves as a natural fish trap when migrating fish seek protection when the water depth is in critical condition, it can also be used as a place to nourish and raise fish in the dry season (Najiyati et al 2005; Bijaksana 2006; Herliwati & Rahman 2011; Sumantriyadi 2014). After the *beje* pond is prepared, then the fish entering the *beje* are allowed to grow naturally without any treatment. At the peak of the dry season the plains around *beje* dry, water and fish in *beje* isolated and trapped, then the fish can be harvested (Rupawan 2004). Types of fish that are harvested are generally black fish groups that have habitat in the swampy marsh, such as common snakehead (*Channa striata*), giant snakehead (*Channa micropeltes*), climbing perch (*Anabas testudineus*), snakeskin gourami (*Trichopodus pectoralis*), three spot gourami (*Trichopodus trichopterus*) and other swamp fish (Burnawi 2009), and these fish belong to important economically valuable fish (Sofia 2017). Some fish species with very high demand, and the specific fish produced only from local wetlands will lead to very high prices in the market, thus providing a significant effect on fisherman's income (Deka et al 2001). This indicates that *beje* has the potential to be further developed, where changes in natural ecosystems are relatively small, even able to maintain local fish species, and support food security and income sources for local communities.

However, the number of units and the size of *beje* managed by local fishermen is based solely on their own capital and experience, so that many wetlands are opened as widely as possible without considering financial feasibility aspects. While Barbier et al (1997) states that for the purpose of wetland conservation, policy is required that does not neglect the loss or degradation of further wetlands through sustainable use and research to measure the value of wetlands. Hence, the objective of the present study was to determine the volume and value of production, as well as the feasibility of *beje* fishery business based on the number of management units.

Material and Method

Description of the study sites. North Hulu Sungai is one of South Kalimantan region with an area of wetlands to $\pm 50,000$ ha. Almost 98.82% of the marsh waters are periodically flooded (CBS North Hulu Sungai Regency 2014) and swamplands will experience drought for some time during the dry season. Drought conditions are used by local communities to trap fish that are trying to find a source of water by making a well dug in the swamp land. This research was conducted in Sungai Papuyu Village Babirik District which is one of *beje* fishery development area in North Hulu Sungai Regency in January-April 2017 (Figure 1). Locations are deliberately chosen based on the number of fisherman households working on *beje* and the variety of fisherman-run units. The *beje* fishery in the study location is managed by 20 fisherman households with total *beje* of 145 units. The number of *beje* managed by each fisherman household varies between 5 and 15 units.

Population and sample. The population in this research is fisherman households who work on *beje* fishery with the number of members as many as 20 households. The members of the population are all sampled. The sample will be divided into three groups based on the number of ownership of *beje* units cultivated. Group I as a small unit is a household that seeks 1 to 5 units; group II as middle unit is households that work on 6 to 10 units; and group III as large unit is households that work on more than 10 units.




Figure 1. The location of *beje* at Babirik district, Hulu Sungai Utara, Indonesia.

Data analysis. The long-term business feasibility of *beje* fishery is determined using the investment criteria approach (Gittinger 1986), as follows:

a. Net Present Value (NPV):

$$NPV = \sum_{t=1}^n \frac{B_t - C_t}{(1+i)^t}$$

where: B_t = total benefit in year- t ;

C_t = total costs in year- t ;

n = the economic life of the project;

i = discount rate.

Decision criteria: $NPV > 0$ - profitable business to be developed further; $NPV = 0$ - business is on break even; $NPV < 0$ - unprofitable business to be further developed.

b. Benefit Cost Ratio (BCR):

$$BCR = \frac{\sum_{t=1}^n B_t}{\sum_{t=1}^n C_t}$$

Decision criteria: $BCR < 1$ - unprofitable business to run; $BCR > 1$ - profitable business to run.

c. Internal Rate of Return (IRR):

$$IRR = i' + \frac{NPV'}{NPV' - NPV} (i - i')$$

where: i_1 = the rate of discount rate that produces NPV_1 (the smallest positive);

i_2 = the rate of discount rate that produces NPV_2 (the smallest negative).

Decision criteria: $IRR < \text{discount rate}$ - unprofitable business to run; $IRR > \text{discount rate}$ - profitable business to run.

Results. *Beje* fisheries have long been worked on by local fishermen, of which 65% of fishermen have been working on it for 5-10 years and 35% of other fishermen have been working on it for more than 10 years. *Beje* which is managed by fishermen has generally a rectangular shape with area based on ownership group is small group around 57.80-462.40 m²; the middle group ranges from 144.50 to 1,300.50 m²; and the average pond size of a large group is 722.50 m² (Table 1). Total production of *beje* per household ranges from 338.64 to 920 kg year⁻¹. Fish produced generally consists of snakeskin gourami (*T. pectoralis*) as much as 103.57-345 kg; three spot gourami (*T. trichopterus*) as much as 147.5-355 kg; common snakehead (*C. striata*) as much as 41.71-115.46 kg; and climbing perch (*A. testudineus*) as much as 45.86-110 kg. While production per unit *beje* ranged from 74 to 100.91 kg year⁻¹ (Table 2).

Table 1

Size of *beje* by unit and household

Group	Size per unit (m ²)			Size per household (m ²)		
	Largest	Smallest	Average	Largest	Smallest	Average
Small	462.40	57.80	310.68	2,312.00	231.20	1,338.69
Middle	1,300.50	144.50	459.77	8,670.00	1,011.50	3,586.23
Large	-	-	722.50	10,837.50	9,392.50	10,115.00

Table 2

Average *beje* production by type of fish and business group (in a year)

Type of fish	Small		Middle		Large	
	Vol. unit ⁻¹ (kg)	Total (kg)	Vol. unit ⁻¹ (kg)	Total (kg)	Vol. unit ⁻¹ (kg)	Total (kg)
Snakeskin gourami	23.71	103.57	28.18	211.82	25	345
Three spot gourami	29.57	147.5	45.00	333.18	27	355
Common snakehead	9.86	41.71	15.00	115.46	8	110
Climbing perch	10.86	45.86	12.73	96.36	8	110
Total	74.00	338.64	100.91	756.82	68	920

The production value of *beje* catch depends on the amount and type of fish produced, as well as the price of the fish. The higher the number and price of the fish caught, the greater the income the fishermen will get. Table 3 shows that snakeskin gourami has the highest production value because its total production is highest, although the selling price is still below the price of common snakehead which is around Rp 15,000-30,000 kg⁻¹. While the second highest production value is common snakehead with a value of Rp 1,131,140-2,875,000; and the selling price per kg reaches Rp 20,000-35,000.

Table 3

Average *beje* production value by type of fish and business group (in a year)

Type of fish	Small		Middle		Large	
	Per unit (Rp 000)	Total (Rp 000)	Per unit (Rp 000)	Total (Rp 000)	Per unit (Rp 000)	Total (Rp 000)
Snakeskin gourami	554.21	2,416.43	430.45	3,243.64	405	5,565
Three spot gourami	147.86	632.14	225.00	1,665.91	135	1,775
Common snakehead	272.00	1,131.14	329.55	2,481.82	205	2,875
Climbing perch	220.43	925.14	259.09	1,940.91	160	2,200
Total	1,194.50	5,104.85	1,244.09	9,332.28	905	12,415

Beje fishery business in each of the average management group requires investment capital of more than Rp 8 million. The highest total investment capital is large group Rp 19,760,000; and the lowest investment is the middle group only Rp 8,755,000. While the operational cost of *beje* in 5 years of highest management is small group Rp 21,577,000; and the lowest operational cost is the middle group Rp 15,428,000. The highest total

revenue and profits are large, while the lowest is the small group (Table 4). Table 4 shows the results of business feasibility analysis with the NPV criterion with a 9% factor discount indicating that in each group can generate profit, where the smallest profit in small group (Rp 3,782,000) and the biggest profit in large group (Rp 44,812,000). The result of NPV analysis with 12% discount factor also still give advantage to each management group. While the result of business feasibility analysis based on BCR criteria at 9% and 12% discount factor, the value of BCR in each group is above 1 which means that each group of management unit is feasible to be cultivated in long term. The lowest BCR value is in the small group and the highest is the large group. While the results of the analysis based on the IRR criteria indicate that the lowest IRR value is small group (16.04%) and the highest is the middle group (19.17%).

Table 4

Beje's business feasibility analysis by business group within 5 years of management

<i>Business feasibility criteria</i>	<i>Small</i>	<i>Middle</i>	<i>Large</i>
Investment cost (Rp 000)	9,918	8,755	19,760
Total cost (Rp 000)	21,577	15,428	15,917
Total revenue (Rp 000)	25,529	46,650	62,150
Profit (Rp 000)	3,952	31,222	46,233
NPV 9% (Rp 000)	3,782	30,280	44,812
NPV 12% (Rp 000)	2,171	21,348	31,387
BCR 9%	1.18	3.00	3.86
BCR 12%	1.13	2.74	3.34
IRR (%)	16.04	19.17	19.01

Discussion. *Beje* is a traditional fishery activity in the inland water swamp where fishing activities are continued with fish polyculture activities. Usually *beje* ponds that have been built by fishermen are left without any treatment until a number of fish from various types of swamp fish are trapped. *Beje* productivity is highly dependent on natural conditions and aquatic fertility, there is no special treatment in management to further encourage fish production, for example an increase in stocking density, feeding, or protection from predators. Therefore, the fish rearing period until harvesting generally takes some time, at least one year. The experience of fishermen shows that the age of *beje* is enough to determine the amount of fish production that can be produced. *Beje* which is old (more than 10 years) allows a lot of fish seeds embedded in it. Adult fish will be harvested immediately, while seeds or young fish are deliberately left and raised to be harvested in the next season.

The results of the study indicate that *beje* business with different size and number of management units can still provide income for fishermen who develop it. The result of business feasibility analysis based on cost and revenue in 5 years of management shows that the three management groups are able to generate profit with ratio between cost and revenue more than 1, so that the three management groups are feasible to be developed further. Similarly, the IRR analysis shows that the IRR of each management group is more than the interest rate for small business loans (12%). Therefore, if the business development uses loan funds with standard bank interest rates, then each level of management still provides benefits for the managers.

However, the highest level of management with productivity per unit was the middle group reaching 100.91 kg year⁻¹ (Table 2). Similarly, the production value per unit of *beje* shows that the economic value of middle group unit (6-10 units) is higher than the other group which is Rp 1,244,090 year⁻¹. While the capital needed for the supply of *beje* units in the middle group is the smallest (Rp 8,755,000). While in large groups (> 10 units) both volume and production value per unit is the lowest at only 68 kg year⁻¹ with production value of Rp 905,000 year⁻¹, but the required total capital is higher is Rp 19,760,000.

Thus, the middle management group has better land use efficiency, capital and operational costs than small and large groups. In line with the results of Sarkar et al

(2015) study, the best annual production level in a dredged pond type culture can be achieved with good management practices, and it comes from small and medium scale cultivation because it is economically more feasible than other scales. Ownership of waters that are not too extensive requires fish farmers to intensify their efforts so that they lead to high productivity (Bairagya 2011). Small-scale polyculture fisheries are feasible (Olawumi et al 2010) with a positive NPV and an IRR of 19-24% (Bigwa 2013). Efforts to increase the productivity of *beje* can be done by developing middle group management unit and more intensifying *beje* maintenance as a fishery cultivation business. Chandra et al (2010) indicate that semi-intensive aquaculture type ponds in floodplain areas are able to provide fish production and high economic benefits for farmers. In addition, in the cultivation of ponds found that the factors that positively affect the production is the extent of the pond area, fish seed, feed, labor and other costs (Tajerin 2007; Onumah & Acquah 2010; Olawumi et al 2010; Adewuyi et al 2010), fish size (Sikiru et al 2009), location and level of water circulation (World Bank 2006), and management capabilities (Ahmed 2007).

Conclusions. The highest total *beje* production volume per unit is the middle group with production reaching 100.91 kg yr⁻¹. The highest production value per unit is in the middle group of Rp 1,244,090 yr⁻¹. The middle management group has better land use, capital and operational efficiency than small and large groups. The results of the study suggest that to increase the productivity of *beje* fisheries can be done by developing *beje* business in the middle group, and supported the application of semi-intensive production technology.

Acknowledgements. We thank to Prof. Dr. Muchlisin Z. A., S.Pi, M.Sc (Dean of Faculty of Marine and Fisheries Syiah Kuala University, Banda Aceh) and two anonymous reviewers for their constructive comments which have helped improve the quality of paper immensely. We are thankful to Fisheries and Husbandry Office of North Hulu Sungai Regency and Fishery Field Extension Officer for information and field assistance for this research, as well as for fishing communities in Babirik swamp area for information and cooperation.

References

- Adewuyi S. A., Phillip B. B., Ayinde I. A., Akerele D., 2010 Analysis of profitability of fish farming in Ogun State, Nigeria. *Journal of Human Ecology* 31(3): 179-184.
- Ahmed N., 2007 Economics of aquaculture feeding practices: Bangladesh. In: Economics of aquaculture feeding practices in selected Asian countries. Hasan M. R. (ed), FAO Fisheries Technical Paper No. 505, Rome, pp. 33-64.
- Bairagya R., 2011 Analysis of profitability of fish farming in West Bengal, India – a study report. *International Journal of Current Research* 3(11):204-214.
- Barbier E. B., Acreman M., Knowler D., 1997 Economic valuation of wetlands: a guide for policy makers and planners. Ramsar Convention Bureau, Gland, Switzerland, 138 pp.
- Bigwa C., 2013 Feasibility of aquaculture in Cameroon: the case of the noun division in the West region [final project]. Fisheries Training Programme, United Nations University, 56 pp.
- Bijaksana U., 2006 [A preliminary study of bio-eco reproduction of snakehead in Bangkau swamp, South Kalimantan Province]. National Symposium of Biotechnology in Aquaculture 2006, Department of Aquaculture Faculty of Fisheries and Marine Sciences, Bogor Agricultural University and Research Institute of Fisheries Freshwater Aquaculture Marine and Fisheries Research Agency, 5 July 2006, 5 pp. [in Indonesian]
- Burnawi B., 2009 [Identification of fish species in fishery sanctuary of Talan Lake, South Kalimantan]. *Buletin Teknik Litkayasa* 7(2):43-46. [in Indonesian]

- CBS (Central Bureau of Statistics) North Hulu Sungai Regency, 2014 [North Hulu Sungai Regency in Figures 2014]. CBS of North Hulu Sungai Regency, South Kalimantan, Indonesia, 493 pp. [in Indonesian]
- Chandra K. J., Sarker D., Khaleque M. A., Das D. R., 2010 Economic analysis of floodplain aquaculture at Daudkandi upazilla in Comilla. *Journal of the Bangladesh Agricultural University* 8(2):323-332.
- Cooke S. J., Allison E. H., Beard Jr. T. D., Arlinghaus R., Arthington A. H., Bartley D. M., Cowx I. G., Fuentesvilla C., Leonard N. J., Lorenzen K., Lynch A. J., Nguyen V. M., Youn S. J., Taylor W. W., Welcomme R. L., 2016 On the sustainability of inland fisheries: finding a future for the forgotten. *Ambio* 45:753-764.
- Das S., Behera B., Mishra A., 2015 Determinants of household use of wetland resources in West Bengal, India. *Wetlands Ecology and Management* 23:803-816.
- David A., Darwall W. R. T., Emerton L., Hamiss H., Hepelwa A., Kasthala G., Kwayu E., Mclvor A., Springate-Baginski O., 2008 An integrated assessment of the biodiversity, livelihood and economic value of wetlands in Mtanza-Msona village, Tanzania. Dar es Salaam, TZ: IUCN Tanzania Country Office, 128 pp.
- Deka T. K., Kakati M., Goswami M. M., 2001 Diversity of wetland fish and its impact on the income of fishermen community of Assam. *Journal of the Indian Fisheries Association* 28:11-22.
- EPA (Environmental Protection Agency), 2006 Economic benefits of wetlands. EPA843-F-06-004, Office of Water, United States of America, 4 pp.
- Früh S., Gattenlöhner U., Hammerl M., Hartmann T., Hörmann S., Megerle H., Spaich F., 2013 Economic value of lakes and wetlands (Summary). Global Nature Fund – University of Applied Forest sciences – Lake Constance Foundation, German, 13 pp.
- Gittinger J. P., 1986 [Economic analysis of agricultural projects]. Second edition, EDI Series in Economic Development, University of Indonesia (UI-Press), Jakarta, 579 pp. [in Indonesian]
- Herliwati, Rahman M., 2011 [Eco-biological characteristics of *beje* fishery in Bangkau swamp area of South Kalimantan]. *Limnotek* 18(1):26-37. [in Indonesian]
- Leschine T. M., Wellman K. F., Green T. H., 1997 The economic value of wetlands: wetlands' role in flood protection in Western Washington. Ecology Publication No. 97-100, National Oceanic and Atmospheric Administration (NOAA), 62 pp.
- Lynch A. J., Cooke S. J., Deines A. M., Bower S. D., Bunnell D. B., Cowx I. G., Nguyen V. M., Nohner J., Phouthavong K., Riley B., Rogers M. W., Taylor W. W., Woelmer W., Youn S. J., Beard Jr. T. D., 2016 The social, economic, and environmental importance of inland fish and fisheries. *Environmental Reviews* 24:115-121.
- Najiyati S., Muslihat L., Suryadiputra I. N. N., 2005 [Guidelines for the management of peatlands for sustainable agriculture]. Climate Change, Forests and Peatlands Projects in Indonesia, Wetlands International - Indonesia Programmed an Wildlife Habitat Canada, Bogor, Indonesia, 198 pp. [in Indonesian]
- Olaoye O. J., Adegbite D. A., Oluwalana E. O., Vaughan I. O., Odebiyi C. O., Adediji A. P., 2014 Comparative evaluation of economic benefits of earthen fish ponds and concrete tanks in aquaculture enterprises in Oyo State, Nigeria. *Croatian Journal of Fisheries* 72:107-117.
- Olawumi A. T., Dipeolu A. O., Bamiro O. M., 2010 Economic analysis of homestead fish production in Ogun State Nigeria. *Journal of Human Ecology* 31(1):13-17.
- Onumah E. E., Acquah H. D., 2010 Frontier analysis of aquaculture farms in the Southern Sector of Ghana. *World Applied Sciences Journal* 9(7):826-835.
- Rupawan, 2004 [*Beje* as a production pond in the swamp land]. Research Center for Inland Water Fisheries Palembang, Indonesia, pp. 189-195. [in Indonesian].
- Sarkar S., Kumar A., Kumar D., Sethi L. N., Hazarika M. M., Das G., 2015 Optimal size of fish pond for socio-economical development of Cachar (Assam). *International Journal of Agriculture, Environment and Biotechnology* 8(2):405-411.
- Sikiru B. O., Omobolanle N. M., Ayorinde B. J. O., Adegoke O. O., 2009 Improving *Clarias* productivity towards achieving food security in Ijebu-Ode, Ogun State, Nigeria: a socioeconomic analysis. *Advances in Biological Research* 3(1-2):24-28.

- Sofia L. A., 2017 Fishermen's perception and participation in fisheries resources conservation of Batungap swamp in Tapin Regency, South Kalimantan, Indonesia. *AAAL Bioflux* 10(6):1618-1626.
- Stuip M. A. M., Baker C. J., Oosterberg W., 2002 The socio-economics of wetlands. Wetlands International and RIZA, The Netherlands, 34 pp.
- Sumantriyadi, 2014 [Utilization of Lebak swamp water resources for fisheries]. *Jurnal Ilmu-ilmu Perikanan dan Budidaya Perairan* 9(1):59-65. [in Indonesian]
- Tajerin, 2007 [Technical efficiency of catfish growth in pond culture (case study in Tulung Agung Regency, East Java Province)]. *Jurnal Ekonomi Pembangunan* 12(1):37-48. [in Indonesian]
- Whiteoak K., Binney J., 2012 Literature review of economic value of ecosystem service that wetlands provide. Masden Jacob Associates, Australia, 70 pp.
- World Bank, 2006 Guidelines for environmental management of aquaculture investments in Vietnam. 236 pp.

Received: 18 october 2018. Accepted: 26 December 2018. Published online: 06 February 2019.

Authors:

Leila Ariyani Sofia, Study Program of Fisheries Agribusiness, Faculty of Fisheries and Marine, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: leila.ariyani@ulm.ac.id

Susi Nurlianti, Graduate of Program of Fisheries Agribusiness, Faculty of Fisheries and Marine, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: susinurlianti066@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Sofia L. A., Nurlianti S., 2019 The economic value of the resource utilization of wetlands: comparative study of *beje* fisheries in North Hulu, Sungai Regency, South Kalimantan, Indonesia. *AAAL Bioflux* 12(1):143-150.